

DADE'S INVARIANT CONJECTURE FOR THE REE GROUPS ${}^2F_4(q^2)$ IN DEFINING CHARACTERISTIC

FRANK HIMSTEDT AND SHIH-CHANG HUANG

ABSTRACT. We verify Dade's invariant conjecture for the simple Ree groups ${}^2F_4(2^{2n+1})$ for all $n > 0$ in the defining characteristic, i.e., in characteristic 2. Together with the results in [3], this completes the proof of Dade's conjecture for the simple Ree groups ${}^2F_4(2^{2n+1})$.

1. INTRODUCTION

The representation theory of finite groups received great impetus from several conjectures linking the representation theory of a finite group G to that of local subgroups, the normalizers of nontrivial p -subgroups of G . A prominent family of such conjectures are Dade's conjectures, published in [9], [10] and [11]. These conjectures express the number of complex irreducible characters with a fixed defect in a given p -block of a finite group G as an alternating sum of related values for p -blocks of certain p -local subgroups.

Dade's conjectures are closely related to several other local-global conjectures: Dade's projective conjecture implies Alperin's weight conjecture, the McKay conjecture and its blockwise version due to Alperin, and one direction of Brauer's height-0 conjecture on abelian blocks. For details see [1], [2], [9], [10], [22], [26] and also [20], [21].

In this paper, we show that Dade's invariant conjecture is true for the simple Ree groups ${}^2F_4(2^{2n+1})$, $n > 0$, in the defining characteristic, i.e., for $p = 2$. Since the Ree groups have a trivial Schur multiplier and a cyclic outer automorphism group for $n > 0$, our results imply that the most general version of Dade's conjectures, the inductive conjecture, is also true for ${}^2F_4(2^{2n+1})$ and $p = 2$. Together with An's results [3] and [4], this completes the proof of Dade's conjectures for the simple groups ${}^2F_4(2^{2n+1})'$ for all $n \geq 0$. Note that this also completes the verification of Uno's invariant conjecture [29] for these groups in the defining characteristic.

The methods we use are similar to those in [16]. By a corollary to the Borel-Tits theorem [6], the normalizers of radical 2-chains are the parabolic subgroups of ${}^2F_4(2^{2n+1})$. Using the character tables of ${}^2F_4(2^{2n+1})$ in the library of the MAPLE [8] part of CHEVIE [14] and the character tables of the parabolic subgroups in [17] and [18], we count characters of the chain normalizers which are fixed by certain outer automorphisms of the Ree groups and then evaluate the alternating sums in Dade's invariant conjecture.

There are some differences between the verification of Dade's conjecture for the Ree groups ${}^2F_4(2^{2n+1})$ and for Steinberg's triality groups ${}^3D_4(2^n)$ in [16]. Firstly, the parabolic subgroups of ${}^2F_4(2^{2n+1})$ have more types of characters which have to

be treated separately. And secondly, when calculating the fixed points of outer automorphisms on the irreducible characters of the parabolic subgroups of ${}^2F_4(2^{2n+1})$ one has to deal with cyclotomic polynomials over $\mathbb{Q}(\sqrt{2})$ instead of $\mathbb{Q}$ making gcd-computations more complicated; see for example Lemma 3.3.

This paper is organized as follows: In Section 2, we fix notation and state Dade's invariant conjecture. In Section 3, we prove some lemmas from elementary number theory which we use to count fixed points of certain automorphisms of ${}^2F_4(2^{2n+1})$. In Section 4, we compute the fixed points of these outer automorphisms on the irreducible characters of parabolic subgroups. In Section 5, we verify Dade's invariant conjecture for ${}^2F_4(2^{2n+1})$ in the defining characteristic. Details on irreducible characters of the Ree groups are summarized in tabular form in Appendix A. In Appendix B, we describe the construction of several families of irreducible characters of ${}^2F_4(2^{2n+1})$. Some of the data in Appendix B might also be of independent interest and is used in [15].

ACKNOWLEDGMENTS

A part of this work was done during visits of the first author at the University of Auckland and at Chiba University. He wishes to express his sincere thanks to all persons of the mathematics departments of these universities for their hospitality, and also to the Marsden Fund of New Zealand and the Japan Society for the Promotion of Science (JSPS) who supported his visits. The second author acknowledges the support of his research during the last years from the Foundation for Research, Science and Technology of New Zealand, the JSPS and the National Science Council, ROC.

2. DADE'S INVARIANT CONJECTURE

Let R be a p -subgroup of a finite group G . Then R is *radical* if $O_p(N(R)) = R$, where $O_p(N(R))$ is the largest normal p -subgroup of the normalizer $N(R) := N_G(R)$. Denote by $\text{Irr}(G)$ the set of all irreducible ordinary characters of G , and by $\text{Blk}(G)$ the set of p -blocks. If $H \leq G$, $\tilde{B} \in \text{Blk}(G)$, and d is an integer, we denote by $\text{Irr}(H, \tilde{B}, d)$ the set of characters $\chi \in \text{Irr}(H)$ satisfying $d(\chi) = d$ and $b(\chi)^G = \tilde{B}$ (in the sense of Brauer), where $d(\chi) = \log_p(|H|_p) - \log_p(\chi(1)_p)$ is the p -defect of χ and $b(\chi)$ is the block of H containing χ .

Given a p -subgroup chain $C : P_0 < P_1 < \cdots < P_n$ of G , define the length $|C| := n$, $C_k : P_0 < P_1 < \cdots < P_k$ and

$$N(C) = N_G(C) := N_G(P_0) \cap N_G(P_1) \cap \cdots \cap N_G(P_n).$$

The chain C is said to be *radical* if it satisfies the following two conditions:

- (a) $P_0 = O_p(G)$ and
- (b) $P_k = O_p(N(C_k))$ for $1 \leq k \leq n$.

Denote by $\mathcal{R} = \mathcal{R}(G)$ the set of all radical p -chains of G .

Suppose $1 \rightarrow G \rightarrow E \rightarrow \overline{E} \rightarrow 1$ is an exact sequence, so that E is an extension of G by $\overline{E}$. Then E acts on $\mathcal{R}$ by conjugation. Given $C \in \mathcal{R}$ and $\psi \in \text{Irr}(N_G(C))$, let $N_E(C, \psi)$ be the stabilizer of (C, ψ) in E , and $N_{\overline{E}}(C, \psi) := N_E(C, \psi)/N_G(C)$. For $\tilde{B} \in \text{Blk}(G)$, an integer $d \geq 0$ and $U \leq \overline{E}$, let $k(N_G(C), \tilde{B}, d, U)$ be the number of characters in the set

$$\text{Irr}(N_G(C), \tilde{B}, d, U) := \{\psi \in \text{Irr}(N_G(C), \tilde{B}, d) \mid N_{\overline{E}}(C, \psi) = U\}.$$

Dade's invariant conjecture can be stated as follows:

Dade's Invariant Conjecture ([11]) *If $O_p(G) = 1$ and $\tilde{B} \in \text{Blk}(G)$ with defect group $D(\tilde{B}) \neq 1$, then*

$$\sum_{C \in \mathcal{R}/G} (-1)^{|C|} k(N_G(C), \tilde{B}, d, U) = 0,$$

where $\mathcal{R}/G$ is a set of representatives for the G -orbits of $\mathcal{R}$.

Let $\text{Aut}(G)$ and $\text{Out}(G)$ be the automorphism and outer automorphism groups of G , respectively. We may suppose $\overline{E} = \text{Out}(G)$. If moreover, $\text{Out}(G)$ is cyclic, then we write

$$k(N_G(C), \tilde{B}, d, |U|) := k(N_G(C), \tilde{B}, d, U).$$

For $G = {}^2F_4(2^{2n+1})$, $n > 0$, $\text{Out}(G)$ is cyclic and the Schur multiplier of G is trivial. So the invariant conjecture for G is equivalent to the inductive conjecture.

3. NOTATION AND LEMMAS FROM ELEMENTARY NUMBER THEORY

From now on, we assume that $p = 2$, n is a positive integer and $q^2 = 2^{2n+1}$. We write ϕ_i for the i -th cyclotomic polynomial in q , for example: $\phi_1 = q - 1$, $\phi_2 = q + 1$, $\phi_3 = q^2 + q + 1$, $\phi_4 = q^2 - 1$, $\phi_6 = q^2 - q + 1$, $\phi_8 = q^4 + 1$, $\phi_{12} = q^4 - q^2 + 1$, $\phi_{24} = q^8 - q^4 + 1$. Furthermore, we set $\phi'_8 := q^2 + \sqrt{2}q + 1$, $\phi''_8 := q^2 - \sqrt{2}q + 1$, $\phi'_{24} = q^4 + \sqrt{2}q^3 + q^2 + \sqrt{2}q + 1$, $\phi''_{24} = q^4 - \sqrt{2}q^3 + q^2 - \sqrt{2}q + 1$. We denote by $\mathbb{N} = \{0, 1, 2, \dots\}$ the set of natural numbers including zero. In the next section, we will use the following lemmas, the first one is given by [5, Lemma 3.1].

Lemma 3.1. *Suppose $m, n \in \mathbb{Z}$ with $m, n > 0$. Then $\gcd(2^m - 1, 2^n - 1) = |2^d - 1|$ where $d := \gcd(m, n)$.*

Lemma 3.2. *Let $f = 2n + 1$ and t be a positive integer with $t \mid f$. Then the following hold.*

- (i) $\gcd(2^t - 1, q^2 - 1) = 2^t - 1$.
- (ii) $\gcd(2^t - 1, q^2 + 1) = 1$.
- (iii) $\gcd(2^t + 1, q^2 - 1) = 1$.
- (iv) $\gcd(2^t + 1, q^2 + 1) = 2^t + 1$.

Proof: (i) is clear by Lemma 3.1.

(ii) Suppose $d = \gcd(2^t - 1, q^2 + 1)$. Since $2^t - 1 \mid q^2 - 1$ by (i), it follows that $d \mid \gcd(q^2 - 1, q^2 + 1) = 1$.

(iv) Since t and f are both odd, we have $2^t + 1 = -((-2)^t - 1)$ and $q^2 + 1 = -((-2)^f - 1)$. So Lemma 3.1 implies $\gcd(2^t + 1, q^2 + 1) = \gcd((-2)^t - 1, (-2)^f - 1) = |(-2)^t - 1| = 2^t + 1$.

(iii) Suppose $d = \gcd(2^t + 1, q^2 - 1)$. Since $2^t + 1 \mid q^2 + 1$ by (iv), it follows that $d \mid \gcd(q^2 - 1, q^2 + 1) = 1$. $\square$

Lemma 3.3. *Let $f = 2n + 1$ and t be a positive integer with $t \mid f$. Then there is a nonnegative integer m such that $f - t = 4tm$ or $f - t = 2t \cdot (2m + 1)$ and for $\varepsilon \in \{1, -1\}$ the following holds:*

(i) If $f - t = 4tm$, then

$$\begin{aligned}\gcd(2^{f-t} - 1, q^2 + \varepsilon\sqrt{2}q + 1) &= 2^t + \varepsilon \cdot (-1)^m \cdot 2^{(t+1)/2} + 1, \\ \gcd(2^{f-t} + 1, q^2 + \varepsilon\sqrt{2}q + 1) &= 1.\end{aligned}$$

(ii) If $f - t = 2t \cdot (2m + 1)$, then

$$\begin{aligned}\gcd(2^{f-t} - 1, q^2 + \varepsilon\sqrt{2}q + 1) &= 1, \\ \gcd(2^{f-t} + 1, q^2 + \varepsilon\sqrt{2}q + 1) &= 2^t + \varepsilon \cdot (-1)^{m+1} \cdot 2^{(t+1)/2} + 1.\end{aligned}$$

Proof: Since $t \mid f = 2n + 1$, the integer t is odd and the even integer $f - t$ is a multiple of $2t$. So, there is $m \in \mathbb{N}$ such that $f - t = 2t \cdot 2m$ or $f - t = 2t \cdot (2m + 1)$.

(i) Suppose $f - t = 4tm$ and $\varepsilon \in \{1, -1\}$. We proceed in several steps:

Step 1: $\gcd(2^{f-t} - 1, q^4 + 1) = 2^{2t} + 1$.

Since $t \mid f$, there is an odd natural number f_u such that $f = t \cdot f_u$. So $f - t = t \cdot (f_u - 1) = 4t \cdot \frac{f_u - 1}{4}$. Hence $2^{f-t} - 1 = (-2^{2t})^{2 \cdot \frac{f_u - 1}{4}} - 1$ and $q^4 + 1 = -((-2^{2t})^{f_u} - 1)$. So Lemma 3.1 implies $\gcd(2^{f-t} - 1, q^4 + 1) = \gcd((-2^{2t})^{2 \cdot \frac{f_u - 1}{4}} - 1, (-2^{2t})^{f_u} - 1) = |(-2^{2t}) - 1| = 2^{2t} + 1$.

Step 2: The number $q^2 + \varepsilon\sqrt{2}q + 1$ is a multiple of $2^t + \varepsilon \cdot (-1)^m \cdot 2^{(t+1)/2} + 1$.

Since $f - t = 2n + 1 - t = 4tm$, we have $n = (4tm + t - 1)/2$. Modulo $2^{2t} + 1$, we compute:

$$\begin{aligned}q^2 + \varepsilon\sqrt{2}q + 1 &\equiv 2^{2n+1} + \varepsilon \cdot 2^{n+1} + 1 = 2^{4tm+t} + \varepsilon \cdot 2^{(4tm+t+1)/2} + 1 \\ &\equiv (2^{2t})^{2m} \cdot 2^t + \varepsilon \cdot (2^{2t})^m \cdot 2^{(t+1)/2} + 1 \\ &\equiv 2^t + \varepsilon \cdot (-1)^m \cdot 2^{(t+1)/2} + 1.\end{aligned}$$

So, since $2^{2t} + 1 = (2^t + \varepsilon \cdot (-1)^m \cdot 2^{(t+1)/2} + 1) \cdot (2^t - \varepsilon \cdot (-1)^m \cdot 2^{(t+1)/2} + 1)$, we can conclude that $q^2 + \varepsilon\sqrt{2}q + 1$ is a multiple of $2^t + \varepsilon \cdot (-1)^m \cdot 2^{(t+1)/2} + 1$.

Step 3: $\gcd(2^{f-t} - 1, q^2 + \varepsilon\sqrt{2}q + 1) = 2^t + \varepsilon \cdot (-1)^m \cdot 2^{(t+1)/2} + 1$.

Since $q^4 + 1 = (q^2 + \sqrt{2}q + 1) \cdot (q^2 - \sqrt{2}q + 1)$ and $\gcd(q^2 + \sqrt{2}q + 1, q^2 - \sqrt{2}q + 1) = 1$, we obtain from Step 1:

$$\begin{aligned}\gcd(2^{f-t} - 1, q^4 + 1) &= \gcd(2^{f-t} - 1, q^2 + \varepsilon\sqrt{2}q + 1) \cdot \gcd(2^{f-t} - 1, q^2 - \varepsilon\sqrt{2}q + 1) \\ &= 2^{2t} + 1 = (2^t + \varepsilon \cdot (-1)^m \cdot 2^{(t+1)/2} + 1) \cdot (2^t - \varepsilon \cdot (-1)^m \cdot 2^{(t+1)/2} + 1).\end{aligned}$$

So, Step 2 implies $\gcd(2^{f-t} - 1, q^2 + \varepsilon\sqrt{2}q + 1) = 2^t + \varepsilon \cdot (-1)^m \cdot 2^{(t+1)/2} + 1$.

Step 4: $\gcd(2^{f-t} + 1, q^2 + \varepsilon\sqrt{2}q + 1) = 1$.

Let $d = \gcd(2^{f-t} + 1, q^4 + 1)$. As in Step 1, we have $f = t \cdot f_u$ and $f - t = t \cdot (f_u - 1) = 4t \cdot \frac{f_u - 1}{4}$ for an odd integer f_u . Since $d \mid 2^{f-t} + 1$ and $d \mid 2^{2f} + 1 = q^4 + 1$, we have $d \mid 2^{2(f-t)} - 1$ and $d \mid 2^{4f} - 1$. By Lemma 3.1, it follows that $\gcd(2^{2(f-t)} - 1, 2^{4f} - 1) = \gcd(2^{4t \cdot 2 \cdot \frac{f_u - 1}{4}} - 1, 2^{4t \cdot f_u} - 1) = 2^{4t} - 1$. Then $d \mid 2^{4t} - 1$, so $2^{4t} \equiv 1 \pmod{d}$ and $0 \equiv 2^{f-t} + 1 = 2^{4t \cdot \frac{f_u - 1}{4}} + 1 \equiv 2 \pmod{d}$. So $d = 1$ as d is odd. Since $q^2 + \varepsilon\sqrt{2}q + 1$ is a divisor of $q^4 + 1$, the claim follows.

(ii) Suppose $f - t = 2t \cdot (2m + 1)$ and $\varepsilon \in \{1, -1\}$. Let $d = \gcd(2^{f-t} - 1, q^4 + 1)$. Then there is an odd $f_u \in \mathbb{N}$ such that $f = t \cdot f_u$, $f - t = t \cdot (f_u - 1) = 2t \cdot \frac{f_u - 1}{2}$. Since $d \mid q^4 + 1 = 2^{2f} + 1$, we have $d \mid 2^{4f} - 1$. By Lemma 3.1, it follows that $\gcd(2^{4f} - 1, 2^{f-t} - 1) = \gcd(2^{2t \cdot 2f_u} - 1, 2^{2t \cdot \frac{f_u - 1}{2}} - 1) = 2^{2t} - 1$. Then $d \mid 2^{2t} - 1$, so $2^{2t} \equiv 1 \pmod{d}$ and $0 \equiv q^4 + 1 = 2^{2f} + 1 = 2^{2t \cdot f_u} + 1 \equiv 2 \pmod{d}$. So $d = 1$ as d is

odd. Since $q^2 + \varepsilon\sqrt{2}q + 1$ is a divisor of $q^4 + 1$, the first statement of (ii) follows. The proof of the second statement of (ii) is analogous to Steps 1-3 of (i). $\square$

4. ACTION OF AUTOMORPHISMS ON IRREDUCIBLE CHARACTERS

Let $G = {}^2F_4(q^2)$ be the simple Ree group with $q^2 = 2^{2n+1}$ and n a positive integer. Let $O = \text{Out}(G)$ and $A = \text{Aut}(G)$. Then $O = \langle \alpha \rangle$ and $A = G \rtimes \langle \alpha \rangle$, where α is a field automorphism of (odd) order $2n + 1$. We fix a Borel subgroup B and maximal parabolic subgroups P_a and P_b of G containing B as in [17] and [18]. In particular, α stabilizes B , P_a and P_b .

In this section, we determine the action of $O = \text{Out}(G)$ on the irreducible characters of the chain normalizers. Our notation for the parameter sets of the irreducible characters of G , B , P_a and P_b is similar to the CHEVIE notation and is given in Table A.1 in Appendix A. For a definition and construction of the irreducible characters $\chi_{42}(k), \dots, \chi_{51}(k, l)$, see Appendix B.

The first column of Table A.1 defines a name for the parameter set which parameterizes those characters which are listed in the second column of the table. The characters of G are numbered according to the character table of ${}^2F_4(q^2)$ in CHEVIE, and for the characters of B , P_a , P_b we use the notation from [17] and [18]. The list of parameters in the third column of Table A.1 in Appendix A is of the form

$$k = 0, \dots, n_1 - 1 \quad \text{or} \quad \begin{array}{l} k = 0, \dots, n_1 - 1 \\ l = 0, \dots, n_2 - 1 \end{array}$$

where the integers n_j are given by polynomials in q with coefficients in $\mathbb{Z}[\sqrt{2}]$. In the first case, the parameter k can be substituted by an element of $\mathbb{Z}$, but two parameters which differ by an element of $n_1\mathbb{Z}$ yield the same character. In the second case, the parameter vector (k, l) can be substituted by an element of $\mathbb{Z} \times \mathbb{Z}$, but two parameter vectors which differ by an element of $n_1\mathbb{Z} \times n_2\mathbb{Z}$ yield the same character. In other words, k can be taken to be an element of $\mathbb{Z}_{n_1}$ and (k, l) can be taken to be an element of $\mathbb{Z}_{n_1} \times \mathbb{Z}_{n_2}$. The groups $\mathbb{Z}_{n_1}$ and $\mathbb{Z}_{n_1} \times \mathbb{Z}_{n_2}$ are also called *character parameter groups* (see Section 3.7 of the CHEVIE [14] manual). The next lines of Table A.1 list elements which have to be excluded from the character parameter group. The remaining parameters are called *admissible* in the following. Different values of admissible parameters may give the same character. The fourth column of Table A.1 defines an equivalence relation on the set of admissible parameters. If no equivalence relation is listed we mean the identity relation. The parameter set is defined to be the set of these equivalence classes. Finally, the last column of Table A.1 gives the cardinality of the parameter set.

We consider the example $P_b I_5$. The character parameter group is $\mathbb{Z}_{q^2-1} \times \mathbb{Z}_{q^2-1}$. The parameter vectors (k, l) and $(k, -l)$ yield the same character and the equivalence class of (k, l) is $\{(k, l), (k, -l)\}$. Hence, the characters $P_b \chi_5(k, l)$ are parameterized by the set

$$P_b I_5 = \{ \{(k, l), (k, -l)\} \mid (k, l) \in \mathbb{Z}_{q^2-1} \times \mathbb{Z}_{q^2-1}, l \neq 0 \}.$$

If we want to emphasize the dependence of a parameter set, say $P_b I_5$, from q we write $P_b I_5(q)$. Table A.1 does not give any detailed information about the parameter sets GI_{34} , GI_{35} , GI_{36} , $\dots$, GI_{41} , GI_{42} , GI_{46} , GI_{50} , GI_{51} , since we will not need an explicit knowledge of these sets (note that the sets GI_1 , GI_{22} , GI_{26} , GI_{28} , GI_{29} , GI_{32} , GI_{34} , GI_{35} , $\dots$, GI_{41} , GI_{42} , GI_{46} , GI_{50} , GI_{51} parameterize the semisimple

irreducible characters of G). The data in Table A.1 is taken from CHEVIE library, the Appendix of [17] and [18] and Appendix B of this paper.

The action of $O = \text{Out}(G)$ on the conjugacy classes of elements of G , B , P_a and P_b induces an action of O on the sets $\text{Irr}(G)$, $\text{Irr}(B)$, $\text{Irr}(P_a)$ and $\text{Irr}(P_b)$ and then an action on the parameter sets. Using the values of the irreducible characters of G , B , P_a and P_b on the classes listed in the last column of Tables A.2-A.15 we can describe the action of O on the parameter sets.

For an O -set I and each subgroup $H \leq O$ let $C_I(H)$ denote the set of fixed points of I under the action of H . In the following proposition we determine $|C_I(H)|$ where I runs through all (disjoint) unions of parameter sets which are listed in Table A.16 except for ${}_GI^{ss}$, where ${}_GI^{ss}$ is the disjoint union of parameter sets parameterizing the semisimple irreducible characters of G . The number of fixed points of ${}_GI^{ss}$ under the action of H was computed in [17, Lemma 6.2].

Proposition 4.1. *Let $t \mid 2n + 1$ and $I \neq {}_GI^{ss}$ be one of the (disjoint) unions of parameter sets listed in Table A.16. If $H = \langle \alpha^t \rangle$ is a subgroup of O , then the second column of Table A.16 shows the number of fixed points $|C_I(H)|$ of I under the action of H .*

Proof: We have to consider the following parameter sets I .

First let

$$\begin{aligned} I \in \{ & {}_GI_2 \cup {}_GI_3, {}_GI_4 \cup {}_GI_{30}, {}_GI_5 \cup {}_GI_6 \cup {}_GI_8 \cup {}_GI_9 \cup {}_GI_{11} \cup {}_GI_{12} \cup {}_GI_{13} \cup {}_GI_{14}, \\ & {}_GI_7 \cup {}_GI_{10}, {}_GI_{17}, {}_GI_{18}, {}_GI_{19} \cup {}_GI_{20}, {}_GI_{31}, {}_BI_6 \cup {}_BI_7, {}_BI_8, {}_BI_{10}, {}_BI_{13} \cup {}_BI_{14}, \\ & {}_BI_{15} \cup {}_BI_{16} \cup {}_BI_{17} \cup {}_BI_{18} \cup {}_BI_{19} \cup {}_BI_{20} \cup {}_BI_{21} \cup {}_BI_{22}, {}_BI_{28} \cup {}_BI_{29}, {}_BI_{43}, \\ & {}_BI_{53} \cup {}_BI_{54}, {}_BI_{55} \cup {}_BI_{56} \cup {}_BI_{57} \cup {}_BI_{58}, {}_PI_6, {}_PI_9 \cup {}_PI_{10}, {}_PI_{11}, {}_PI_{13} \cup {}_PI_{25}, \\ & {}_PI_{14} \cup {}_PI_{15} \cup {}_PI_{16} \cup {}_PI_{17} \cup {}_PI_{21} \cup {}_PI_{22} \cup {}_PI_{23} \cup {}_PI_{24}, {}_PI_{20}, {}_PI_{29}, {}_PI_{30}, \\ & {}_PI_{31}, {}_PI_9 \cup {}_PI_{10}, {}_PI_{11}, {}_PI_{13}, {}_PI_{14} \cup {}_PI_{15}, {}_PI_{17} \cup {}_PI_{18} \cup {}_PI_{19} \cup {}_PI_{20} \\ & \cup {}_PI_{22} \cup {}_PI_{23} \cup {}_PI_{24} \cup {}_PI_{25}, {}_PI_{29} \cup {}_PI_{30}, {}_PI_{43} \cup {}_PI_{50}, {}_PI_{44} \cup {}_PI_{45} \\ & \cup {}_PI_{51} \cup {}_PI_{52}, {}_PI_{46} \cup {}_PI_{53} \}. \end{aligned}$$

The degrees and character values on the conjugacy classes listed in Tables A.2-A.15 show $C_I(H) = I$ and hence $|C_I(H)| = |I|$. We demonstrate this for the parameter set $I = {}_PI_{43} \cup {}_PI_{50}$. The degrees in Table A.5 show that ${}_PI_{43}\chi_{43}$ and ${}_PI_{50}\chi_{50}$ are the only irreducible characters of P_b of degree $\frac{q^9}{\sqrt{2}}(q^2 - 1)$. Hence, ${}_PI_{43}\chi_{43}^\alpha \in \{{}_PI_{43}\chi_{43}, {}_PI_{50}\chi_{50}\}$. The class representatives in Table A.7 in [18] show that the conjugacy class $c_{1,9}$ is fixed by α and we can see from the character table A.9 of P_b in [18] that the values of ${}_PI_{43}\chi_{43}$ and ${}_PI_{50}\chi_{50}$ on $c_{1,9}$ are different. So, ${}_PI_{43}\chi_{43}^\alpha = {}_PI_{43}\chi_{43}$ for $i = 43, 50$ and $|C_I(H)| = |I|$.

In each of the following cases, we have that the action of α on I is given by $x^\alpha = 2x$ for all $x \in I$ using the character values on the classes listed in the last column of Tables A.5-A.15. We demonstrate this for the parameter set $I = {}_PI_3 \cup {}_PI_4$. The degrees in Table A.15 show that the ${}_PI_3\chi_3(k, l)$'s are the only irreducible characters of P_a of degree $q^2 + 1$, so ${}_PI_3\chi_3(k, l)^\alpha = {}_PI_3\chi_3(k', l')$ for some $\{(k', l'), \dots\} \in {}_PI_3$. We see from the class representatives in [18] that α acts on the semisimple conjugacy classes of P_a like the 2nd power map which implies that the values of ${}_PI_3\chi_3(k', l')$ and ${}_PI_3\chi_3(2k, 2l)$ on the semisimple classes coincide. Then, the character values of ${}_PI_3\chi_3(k, l)$ (see the character table A.5 in [18]) imply that the values of ${}_PI_3\chi_3(k', l')$ and ${}_PI_3\chi_3(2k, 2l)$ coincide on all classes, hence ${}_PI_3\chi_3(k', l') = {}_PI_3\chi_3(2k, 2l)$ and therefore ${}_PI_3\chi_3(k, l)^\alpha = {}_PI_3\chi_3(2k, 2l)$. Similarly, ${}_PI_4\chi_4(k)^\alpha = {}_PI_4\chi_4(2k)$. Hence, $x^\alpha = 2x$ for all $x \in I$.

Let $I = {}_G I_{27} \cup {}_G I_{33}$. If $x = \{k, -k\} \in I$, then $x \in C_I(H)$ if and only if $(2^t - 1)k \equiv 0$ or $(2^t + 1)k \equiv 0$. Let $C_{\pm} := \{\{k, -k\} \in C_I(H) \mid (2^t \pm 1)k \equiv 0\}$, so that $C_I(H) = C_- \cup C_+$ and $C_- \cap C_+ = \emptyset$. We claim

$$C_- = \left\{ \{k, -k\} \in {}_G I_{27} \mid k \text{ is a multiple of } \frac{q^2 - 1}{2^t - 1} \right\}.$$

The inclusion $\supseteq$ is clear. Let $x = \{k, -k\} \in C_-$. If $x \in {}_G I_{33}$, then $(2^t - 1)k \equiv 0 \pmod{q^2 + 1}$ and Lemma 3.2(ii) implies $k \equiv 0$, which is impossible. Hence $x \in {}_G I_{27}$ and $(2^t - 1)k \equiv 0 \pmod{q^2 - 1}$. By Lemma 3.2(i), k is a multiple of $(q^2 - 1)/(2^t - 1)$, proving the claim. Next, we claim

$$C_+ = \left\{ \{k, -k\} \in {}_G I_{33} \mid k \text{ is a multiple of } \frac{q^2 + 1}{2^t + 1} \right\}.$$

The inclusion $\supseteq$ is clear. Let $x = \{k, -k\} \in C_+$. If $x \in {}_G I_{27}$, then $(2^t + 1)k \equiv 0 \pmod{q^2 - 1}$ and Lemma 3.2(iii) implies $k \equiv 0$, which is impossible. Hence $x \in {}_G I_{33}$ and $(2^t + 1)k \equiv 0 \pmod{q^2 + 1}$. By Lemma 3.2(iv), k is a multiple of $(q^2 + 1)/(2^t + 1)$ and the claim holds.

Thus in all cases, $|C_I(H)| = |C_-| + |C_+| = \frac{2^t - 2}{2} + \frac{2^t - 2}{2} = 2^t - 2$.

Let $I = {}_B I_1$. If $(k, l) \in I$, then $(k, l) \in C_I(H)$ if and only if $(2^t - 1)k \equiv 0$ and $(2^t - 1)l \equiv 0 \pmod{q^2 - 1}$. By Lemma 3.2(i), this is equivalent with k, l are multiples of $\frac{q^2 - 1}{2^t - 1}$. Thus, $|C_I(H)| = (2^t - 1)^2$.

Now, let I be one of the parameter sets ${}_B I_2, {}_B I_3, {}_B I_4, {}_B I_5, {}_B I_9, {}_B I_{11}, {}_B I_{12}, {}_B I_{23}, {}_B I_{26}, {}_B I_{27}, {}_B I_{39}, {}_B I_{42}, {}_B I_{44}, {}_B I_{45}, {}_B I_{51}, {}_B I_{52}, {}_{P_a} I_1, {}_{P_a} I_2, {}_{P_a} I_5, {}_{P_a} I_7, {}_{P_a} I_8, {}_{P_a} I_{12}, {}_{P_a} I_{26}, {}_{P_a} I_{34}, {}_{P_a} I_{35}, {}_{P_b} I_1, {}_{P_b} I_2, {}_{P_b} I_3, {}_{P_b} I_4, {}_{P_b} I_8, {}_{P_b} I_{12}, {}_{P_b} I_{27}, {}_{P_b} I_{28}, {}_{P_b} I_{40}$.

If $k \in I$, then $k \in C_I(H)$ if and only if $(2^t - 1)k \equiv 0 \pmod{q^2 - 1}$. By Lemma 3.2(i), we get $C_I(H) = \{k \in I \mid k \text{ is a multiple of } (q^2 - 1)/(2^t - 1)\}$ and $|C_I(H)| = 2^t - 1$.

Let $I = {}_{P_a} I_3 \cup {}_{P_a} I_4$. First, we compute $|C_{{}_{P_a} I_3}(H)|$. Let

$$U_i := \begin{cases} \{\{(k, l), (k', l')\} \in C_{{}_{P_a} I_3}(H) \mid 2^t k \equiv k, 2^t l \equiv l\} & \text{if } i = 1, \\ \{\{(k, l), (k', l')\} \in C_{{}_{P_a} I_3}(H) \mid 2^t k \equiv k', 2^t l \equiv l'\} & \text{if } i = 2, \end{cases}$$

where $k' = \frac{q}{\sqrt{2}}(k + l)$ and $l' = \frac{q}{\sqrt{2}}(k - l)$.

If $x = \{(k, l), (\frac{q}{\sqrt{2}}(k + l), \frac{q}{\sqrt{2}}(k - l))\} \in {}_{P_a} I_3$, then $x \in U_1$ if and only if $(2^t - 1)k \equiv 0$ and $(2^t - 1)l \equiv 0 \pmod{q^2 - 1}$. By Lemma 3.2(i), this is equivalent with k, l are multiples of $\frac{q^2 - 1}{2^t - 1}$. Thus, $|U_1| = (2^t - 2)(2^t - 1)/2$.

If $x = \{(k, l), (\frac{q}{\sqrt{2}}(k + l), \frac{q}{\sqrt{2}}(k - l))\} \in {}_{P_a} I_3$, then $x \in U_2$ if and only if $2^t k \equiv \frac{q}{\sqrt{2}}(k + l)$ and $2^t l \equiv \frac{q}{\sqrt{2}}(k - l) \pmod{q^2 - 1}$. Multiplying both congruences by $\sqrt{2}q$, we get $\sqrt{2}q \cdot 2^t k \equiv k + l$ and $\sqrt{2}q \cdot 2^t l \equiv k - l \pmod{q^2 - 1}$. Then, $l \equiv (\sqrt{2}q \cdot 2^t - 1)k$ and $k \equiv (\sqrt{2}q \cdot 2^t + 1)l \pmod{q^2 - 1}$. Combining these two congruences, we get $k \equiv (\sqrt{2}q \cdot 2^t - 1)(\sqrt{2}q \cdot 2^t + 1)k = (2q^2 \cdot 2^{2t} - 1)k \equiv (2 \cdot 2^{2t} - 1)k$. Then $2(2^{2t} - 1)k \equiv 0 \pmod{q^2 - 1}$. By Lemma 3.2(i) and (iii), it follows that $\gcd(2(2^t - 1)(2^t + 1), q^2 - 1) = \gcd(2^t - 1, q^2 - 1) = 2^t - 1$. Thus $(2^t - 1)k \equiv 0 \pmod{q^2 - 1}$. Then $k \equiv 2^t k \equiv \frac{q}{\sqrt{2}}(k + l)$. Multiplying both sides by $\sqrt{2}q$, we get $\sqrt{2}qk \equiv k + l$. But then $l \equiv (\sqrt{2}q - 1)k$, a contradiction to the definition of ${}_{P_a} I_3$. Hence, $U_2 = \emptyset$. So $|C_{{}_{P_a} I_3}(H)| = |U_1| + |U_2| = (2^t - 2)(2^t - 1)/2$.

Next, we calculate $|C_{P_a I_4}(H)|$. If $x = \{k, q^2 k\} \in P_a I_4$, then $x \in C_{P_a I_4}(H)$ if and only if $(2^t - 1)k \equiv 0$ or $(2^t - q^2)k \equiv 0 \pmod{(q^2 + 1)(q^2 - 1)}$. Suppose $(2^t - 1)k \equiv 0$. By Lemma 3.2(i) and (ii), it follows that $\gcd(2^t - 1, (q^2 + 1)(q^2 - 1)) = \gcd(2^t - 1, q^2 - 1) = 2^t - 1$. Thus, $(q^2 + 1) \cdot \frac{(q^2 - 1)}{2^t - 1} \mid k$. But then $(q^2 + 1) \mid k$, a contradiction to the definition of $P_a I_4$. So we have proved that $x \in C_{P_a I_4}(H)$ if and only if $(2^t - q^2)k \equiv 0 \pmod{(q^2 + 1)(q^2 - 1)}$. We claim

$$C_{P_a I_4}(H) = \left\{ \{k, q^2 k\} \in P_a I_4 \mid k \text{ is a multiple of } \frac{(q^2 + 1)(q^2 - 1)}{(2^t + 1)(2^t - 1)} \right\}.$$

Let $k = \frac{(q^2 + 1)(q^2 - 1)}{(2^t + 1)(2^t - 1)} \cdot m$ for some $m \in \mathbb{Z}$. Because $t \mid 2n + 1$ we have $2t \mid 2n + 1 - t$. Then we get $(2^t + 1)(2^t - 1) \mid 2^{2n+1-t} - 1$. Thus $(2^{2n+1-t} - 1)k = \frac{2^{2n+1-t} - 1}{(2^t + 1)(2^t - 1)}(q^2 + 1)(q^2 - 1) \cdot m \equiv 0 \pmod{(q^2 + 1)(q^2 - 1)}$. So $(2^t - q^2)k \equiv 0 \pmod{(q^2 + 1)(q^2 - 1)}$ and $\{k, q^2 k\} \in C_{P_a I_4}(H)$.

Conversely, suppose $\{k, q^2 k\} \in C_{P_a I_4}(H)$. Then $(2^t - q^2)k \equiv 0 \pmod{(q^2 + 1)(q^2 - 1)}$. Hence $(2^t + 1)k \equiv 0 \pmod{q^2 + 1}$ and $(2^t - 1)k \equiv 0 \pmod{q^2 - 1}$. By Lemma 3.2(i) and (iv), this is equivalent with $\frac{q^2 + 1}{2^t + 1} \mid k$ and $\frac{q^2 - 1}{2^t - 1} \mid k$. Since $\frac{q^2 + 1}{2^t + 1} \mid q^2 + 1$ and $\frac{q^2 - 1}{2^t - 1} \mid q^2 - 1$ and since $\gcd(q^2 + 1, q^2 - 1) = 1$, we have $\gcd\left(\frac{q^2 + 1}{2^t + 1}, \frac{q^2 - 1}{2^t - 1}\right) = 1$. Therefore $\frac{(q^2 + 1)(q^2 - 1)}{(2^t + 1)(2^t - 1)} \mid k$, and the claim holds. So by the definition of $P_a I_4$, we get $|C_{P_a I_4}(H)| = 2^t(2^t - 1)/2$.

So, $|C_I(H)| = |C_{P_a I_3}(H)| + |C_{P_a I_4}(H)| = (2^t - 1)^2$.

Let $I = P_a I_{32} \cup P_a I_{33}$. The computation of $|C_I(H)|$ is analogous to the case $I = G I_{27} \cup G I_{33}$. One gets $|C_I(H)| = 2^t - 1$.

Let $I = P_b I_5 \cup P_b I_6 \cup P_b I_7$. First, we compute $|C_{P_b I_5}(H)|$. Let

$$U_i := \begin{cases} \left\{ \{(k, l), (k, -l)\} \in C_{P_b I_5}(H) \mid 2^t k \equiv k, 2^t l \equiv l \right\} & \text{if } i = 1, \\ \left\{ \{(k, l), (k, -l)\} \in C_{P_b I_5}(H) \mid 2^t k \equiv k, 2^t l \equiv -l \right\} & \text{if } i = 2. \end{cases}$$

If $x = \{(k, l), (k, -l)\} \in P_b I_5$, then $x \in U_1$ if and only if $(2^t - 1)k \equiv 0$ and $(2^t - 1)l \equiv 0 \pmod{q^2 - 1}$. By Lemma 3.2(i), this is equivalent with k, l are multiples of $\frac{q^2 - 1}{2^t - 1}$. Thus, $|U_1| = (2^t - 1)(2^t - 2)/2$.

If $x = \{(k, l), (k, -l)\} \in P_b I_5$, then $x \in U_2$ if and only if $(2^t - 1)k \equiv 0$ and $(2^t + 1)l \equiv 0 \pmod{q^2 - 1}$. By Lemma 3.2(iii), the second congruence is equivalent with $l \equiv 0 \pmod{q^2 - 1}$, a contradiction to the definition of $P_b I_5$. Hence $U_2 = \emptyset$. So $|C_{P_b I_5}(H)| = |U_1| + |U_2| = (2^t - 1)(2^t - 2)/2$.

Next, we calculate $|C_{P_b I_6 \cup P_b I_7}(H)|$. Let $\phi'_8 := q^2 + \sqrt{2}q + 1$ and $\phi''_8 := q^2 - \sqrt{2}q + 1$. The set $J_1 := \left\{ \{(k, l), (k, -l), (k, q^2 l), (k, -q^2 l)\} \mid (k, l) \in \mathbb{Z}_{q^2 - 1} \times \mathbb{Z}_{q^2 + \sqrt{2}q + 1}, l \neq 0 \right\}$ becomes an H -set via $x^\alpha := 2x$ for all $x \in J$, and we have $J_1 \simeq P_b I_6$ as H -sets; note that $P_b \chi_6(k)$ is obtained via inflation from a regular semisimple character of the Levi complement $L_b \cong \mathbb{Z}_{q^2 - 1} \times \text{Sz}(q^2)$. In a similar way, we have an isomorphism of H -sets

$$P_b I_7 \simeq \left\{ \{(k, l), (k, -l), (k, q^2 l), (k, -q^2 l)\} \mid (k, l) \in \mathbb{Z}_{q^2 - 1} \times \mathbb{Z}_{q^2 - \sqrt{2}q + 1}, l \neq 0 \right\} =: J_2.$$

Hence, ${}_{P_b}I_6 \cup {}_{P_b}I_7 \simeq J_1 \cup J_2$ (disjoint union) as H -sets, so that we can identify ${}_{P_b}I_6 \cup {}_{P_b}I_7 = J_1 \cup J_2$.

In a first step, we compute $|C_{J_1}(H)|$. Let

$$\begin{aligned} U_{\pm 1} &:= \{(k, l), (k, -l), (k, q^2l), (k, -q^2l)\} \in C_{J_1}(H) \mid 2^t k \equiv k, 2^t l \equiv \pm l\}, \\ U_{\pm 2} &:= \{(k, l), (k, -l), (k, q^2l), (k, -q^2l)\} \in C_{J_1}(H) \mid 2^t k \equiv k, 2^t l \equiv \pm q^2l\}, \end{aligned}$$

where the congruences are mod $q^2 - 1$ and $q^2 + \sqrt{2}q + 1$, respectively. Then $C_{J_1}(H) = U_1 \cup U_{-1} \cup U_2 \cup U_{-2}$. We claim that $U_{\pm 1} = \emptyset$. Suppose $\{(k, l), \dots\} \in U_1$. Then $(2^t - 1)k \equiv 0 \pmod{q^2 - 1}$ and $(2^t - 1)l \equiv 0 \pmod{q^2 + \sqrt{2}q + 1}$. Since $2^t - 1 \mid q^2 - 1 \mid q^4 - 1$ and since $\gcd(q^4 - 1, (q^2 + \sqrt{2}q + 1) \cdot (q^2 - \sqrt{2}q + 1)) = \gcd(q^4 - 1, q^4 + 1) = 1$, the second congruence is equivalent with $l \equiv 0 \pmod{q^2 + \sqrt{2}q + 1}$, contradicting the definition of J_1 . So $U_1 = \emptyset$. Suppose $\{(k, l), \dots\} \in U_{-1}$. Then $(2^t - 1)k \equiv 0 \pmod{q^2 - 1}$ and $(2^t + 1)l \equiv 0 \pmod{q^2 + \sqrt{2}q + 1}$. Since $2^t + 1 \mid q^2 + 1 \mid q^4 - 1$ and since $\gcd(q^4 - 1, q^4 + 1) = 1$, the second congruence is equivalent with $l \equiv 0 \pmod{q^2 + \sqrt{2}q + 1}$, contradicting the definition of J_1 . So $U_{-1} = \emptyset$.

Suppose $\{(k, l), \dots\} \in U_2$. Then $2^t k \equiv k \pmod{q^2 - 1}$ and $2^t l \equiv q^2 l \pmod{q^2 + \sqrt{2}q + 1}$, and hence $(2^t - 1)k \equiv 0 \pmod{q^2 - 1}$ and $(2^{2n+1-t} - 1)l \equiv 0 \pmod{q^2 + \sqrt{2}q + 1}$. By Lemmas 3.2(i), 3.3 and the definition of J_1 , we get

$$|U_2| = \begin{cases} \frac{1}{4}(2^t - 1)(2^t \pm 2^{(t+1)/2}) & \text{if } 4t \mid 2n + 1 - t, \\ 0 & \text{if } 4t \nmid 2n + 1 - t. \end{cases}$$

Suppose $\{(k, l), (k, -l), (k, q^2l), (k, -q^2l)\} \in U_{-2}$. Then $2^t k \equiv k \pmod{q^2 - 1}$ and $2^t l \equiv -q^2 l \pmod{q^2 + \sqrt{2}q + 1}$, and hence $(2^t - 1)k \equiv 0 \pmod{q^2 - 1}$ and $(2^{2n+1-t} + 1)l \equiv 0 \pmod{q^2 + \sqrt{2}q + 1}$. By Lemma 3.2(i), 3.3 and the definition of J_1 , we get

$$|U_{-2}| = \begin{cases} 0 & \text{if } 4t \mid 2n + 1 - t, \\ \frac{1}{4}(2^t - 1)(2^t \pm 2^{(t+1)/2}) & \text{if } 4t \nmid 2n + 1 - t. \end{cases}$$

So

$$|C_{J_1}(H)| = |U_1| + |U_{-1}| + |U_2| + |U_{-2}| = \frac{1}{4}(2^t - 1)(2^t \pm 2^{(t+1)/2}).$$

Analogously, we obtain

$$|C_{J_2}(H)| = \frac{1}{4}(2^t - 1)(2^t \mp 2^{(t+1)/2}).$$

So

$$|C_{{}_{P_b}I_6 \cup {}_{P_b}I_7}(H)| = |C_{J_1 \cup J_2}(H)| = |C_{J_1}(H)| + |C_{J_2}(H)| = 2^t(2^t - 1)/2.$$

Hence we get

$$|C_I(H)| = |C_{{}_{P_b}I_5}(H)| + |C_{{}_{P_b}I_6 \cup {}_{P_b}I_7}(H)| = \frac{(2^t - 1)(2^t - 2)}{2} + \frac{2^t(2^t - 1)}{2} = (2^t - 1)^2.$$

Let $I \in \{ {}_G I_{23} \cup {}_G I_{43} \cup {}_G I_{47}, {}_G I_{24} \cup {}_G I_{44} \cup {}_G I_{48}, {}_G I_{25} \cup {}_G I_{45} \cup {}_G I_{49}, {}_{P_b} I_{16} \cup {}_{P_b} I_{21} \cup {}_{P_b} I_{26}, {}_{P_b} I_{47} \cup {}_{P_b} I_{48} \cup {}_{P_b} I_{49}, {}_{P_b} I_{54} \cup {}_{P_b} I_{55} \cup {}_{P_b} I_{56} \}$. Then these (disjoint) unions of parameter sets are isomorphic H -sets, so that we can assume $I = {}_{P_b} I_{47} \cup {}_{P_b} I_{48} \cup {}_{P_b} I_{49}$. First, we compute $|C_{{}_{P_b} I_{47}}(H)|$. If $x = \{k, -k\} \in I$, then $x \in C_{{}_{P_b} I_{47}}(H)$ if and only if $(2^t - 1)k \equiv 0$ or $(2^t + 1)k \equiv 0 \pmod{q^2 - 1}$. Let

$$C_{\pm} := \{ \{k, -k\} \in C_{{}_{P_b} I_{47}}(H) \mid (2^t \pm 1)k \equiv 0 \},$$

so that $C_I(H) = C_- \cup C_+$ and $C_- \cap C_+ = \emptyset$. We claim that $C_+ = \emptyset$. Let $x = \{k, -k\} \in C_+$, then $(2^t + 1)k \equiv 0 \pmod{q^2 - 1}$ and Lemma 3.2(iii) implies $k \equiv 0$, which is impossible. Hence $x \in C_-$ and $(2^t - 1)k \equiv 0 \pmod{q^2 - 1}$. By Lemma 3.2(i), k is a multiple of $(q^2 - 1)/(2^t - 1)$ and $|C_I(H)| = |C_-| + |C_+| = \frac{2^t - 2}{2}$.

Next, we compute $|C_{P_b I_{48} \cup P_b I_{49}}(H)|$. We start with $P_b I_{48}$. Let

$$\begin{aligned} U_{\pm 1} &:= \left\{ \{k, -k, q^2 k, -q^2 k\} \in C_{P_b I_{48}}(H) \mid 2^t k \equiv \pm k \right\}, \\ U_{\pm 2} &:= \left\{ \{k, -k, q^2 k, -q^2 k\} \in C_{P_b I_{48}}(H) \mid 2^t k \equiv \pm q^2 k \right\}, \end{aligned}$$

where the congruences are mod $q^2 + \sqrt{2}q + 1$. Then $C_{P_b I_{48}}(H) = U_1 \cup U_{-1} \cup U_2 \cup U_{-2}$. We claim that $U_{\pm 1} = \emptyset$. Suppose $\{k, -k, q^2 k, -q^2 k\} \in U_1$. Then $(2^t - 1)k \equiv 0$. Since $2^t - 1 \mid q^2 - 1 \mid q^4 - 1$ and since $\gcd(q^4 - 1, (q^2 + \sqrt{2}q + 1) \cdot (q^2 - \sqrt{2}q + 1)) = \gcd(q^4 - 1, q^4 + 1) = 1$, we have $k \equiv 0 \pmod{q^2 + \sqrt{2}q + 1}$, contradicting the definition of $P_b I_{48}$. So $U_1 = \emptyset$. Suppose $\{k, -k, q^2 k, -q^2 k\} \in U_{-1}$. Then $(2^t + 1)k \equiv 0 \pmod{q^2 + \sqrt{2}q + 1}$. Since $2^t + 1 \mid q^2 + 1 \mid q^4 - 1$ and since $\gcd(q^4 - 1, q^4 + 1) = 1$, it follows that $k \equiv 0 \pmod{q^2 + \sqrt{2}q + 1}$, contradicting the definition of $P_b I_{48}$. So $U_{-1} = \emptyset$. Suppose $\{k, -k, q^2 k, -q^2 k\} \in U_2$. Then $2^t k \equiv q^2 k \pmod{q^2 + \sqrt{2}q + 1}$ and hence $(2^{2n+1-t} - 1)k \equiv 0 \pmod{q^2 + \sqrt{2}q + 1}$. By Lemma 3.3 and the definition of $P_b I_{48}$, we get

$$|U_2| = \begin{cases} \frac{1}{4}(2^t \pm 2^{(t+1)/2}) & \text{if } 4t \mid 2n + 1 - t, \\ 0 & \text{if } 4t \nmid 2n + 1 - t. \end{cases}$$

Suppose $\{k, -k, q^2 k, -q^2 k\} \in U_{-2}$. Then $2^t k \equiv -q^2 k \pmod{q^2 + \sqrt{2}q + 1}$ and hence $(2^{2n+1-t} + 1)k \equiv 0 \pmod{q^2 + \sqrt{2}q + 1}$. By Lemma 3.3 and the definition of $P_b I_{48}$, we get

$$|U_{-2}| = \begin{cases} 0 & \text{if } 4t \mid 2n + 1 - t, \\ \frac{1}{4}(2^t \pm 2^{(t+1)/2}) & \text{if } 4t \nmid 2n + 1 - t. \end{cases}$$

So

$$|C_{P_b I_{48}}(H)| = |U_1| + |U_{-1}| + |U_2| + |U_{-2}| = \frac{1}{4}(2^t \pm 2^{(t+1)/2}).$$

Analogously, we obtain

$$|C_{P_b I_{49}}(H)| = \frac{1}{4}(2^t \mp 2^{(t+1)/2}).$$

So we have $|C_{P_b I_{48} \cup P_b I_{49}}(H)| = |C_{P_b I_{48}}(H)| + |C_{P_b I_{49}}(H)| = 2^t/2$. Hence, we get $|C_I(H)| = |C_{P_b I_{47}}(H)| + |C_{P_b I_{48} \cup P_b I_{49}}(H)| = \frac{2^t - 2}{2} + \frac{2^t}{2} = 2^t - 1$. $\square$

Proposition 4.2. *Let $t \mid 2n+1$, $H = \langle \alpha^t \rangle \leq O$ and let $(I, J) \in \{(GI_{15} \cup GI_{16}, P_a I_{18} \cup P_a I_{19}), (BI_{24} \cup BI_{25}, P_a I_{27} \cup P_a I_{28}), (BI_{30} \cup BI_{31} \cup BI_{32} \cup BI_{33} \cup BI_{35} \cup BI_{36}, P_b I_{31} \cup P_b I_{32} \cup P_b I_{33} \cup P_b I_{34} \cup P_b I_{36} \cup P_b I_{37}), (BI_{34} \cup BI_{37}, P_b I_{35} \cup P_b I_{38}), (BI_{38} \cup BI_{40} \cup BI_{41}, P_b I_{39} \cup P_b I_{41} \cup P_b I_{42}), (BI_{46} \cup BI_{47} \cup BI_{48} \cup BI_{49}, P_a I_{36} \cup P_a I_{37} \cup P_a I_{38} \cup P_a I_{39}), (BI_{50}, P_a I_{40})\}$. Then $|C_I(H)| = |C_J(H)|$.*

Proof: Suppose $(I, J) = (GI_{15} \cup GI_{16}, P_a I_{18} \cup P_a I_{19})$. Considering degrees we see that the sets $\{G\chi_{15}, G\chi_{16}\}$ and $\{P_a \chi_{18}, P_a \chi_{19}\}$ are invariant under the action of α (note that $P_a \chi_{20}$ is real-valued and $P_a \chi_{18}, P_a \chi_{19}$ are not). The characters $G\chi_{15}, G\chi_{16}$ coincide on all classes of G except for the classes $c_{5,3}$ and $c_{5,4}$. Similarly, $P_a \chi_{18}, P_a \chi_{19}$ coincide on all conjugacy classes of P_a except for $c_{8,3}$ and $c_{8,4}$. From

the class representatives in [17, Table A.2] and [18, Table A.3] we get that α swaps ${}_G\chi_{15}$ and ${}_G\chi_{16}$ if and only if it swaps ${}_{P_a}\chi_{18}$ and ${}_{P_a}\chi_{19}$. So, $|C_I(H)| = |C_J(H)|$.

Next, suppose $(I, J) = ({}_B I_{50}, {}_{P_a} I_{40})$. By construction, ${}_{P_a}\chi_{40}(k) = {}_B\chi_{50}(k)^{P_a}$ is induced from the α -stable Borel subgroup B for all $k \in {}_B I_{50} = {}_{P_a} I_{40}$ (see [18, Table 3]) and induction of characters induces a bijection from $\{{}_B\chi_{50}(k) \mid k \in {}_B I_{50}\}$ onto $\{{}_{P_a}\chi_{40}(k) \mid k \in {}_{P_a} I_{40}\}$ mapping ${}_B\chi_{50}(k)$ to ${}_{P_a}\chi_{40}(k)$. We have

$${}_{P_a}\chi_{40}(k^\alpha) = {}_{P_a}\chi_{40}(k)^\alpha = ({}_B\chi_{50}(k)^{P_a})^\alpha = ({}_B\chi_{50}(k)^\alpha)^{P_a} = ({}_B\chi_{50}(k^\alpha))^{P_a}.$$

So the above-mentioned bijection is an isomorphism of H -sets. Hence, ${}_B I_{50} \simeq {}_{P_a} I_{40}$ as H -sets and $|C_{{}_B I_{50}}(H)| = |C_{{}_{P_a} I_{40}}(H)|$. The remaining pairs of parameter sets can be treated analogously, see the construction of the characters ${}_{P_a}\chi_{28}(k)$, ${}_{P_a}\chi_{38}(k)$, ${}_{P_a}\chi_{39}(k)$, ${}_{P_b}\chi_{36}(k)$, ${}_{P_b}\chi_{37}(k)$, ${}_{P_b}\chi_{38}(k)$ and ${}_{P_b}\chi_{42}(k)$ in [18]. $\square$

5. DADE'S INVARIANT CONJECTURE FOR ${}^2F_4(q^2)$

In this section, we prove Dade's invariant conjecture for $G = {}^2F_4(2^{2n+1})$, $n > 0$, in the defining characteristic. As in the previous section, let $O = \text{Out}(G) = \langle \alpha \rangle$, where α is a field automorphism of (odd) order $2n + 1$. We fix a Borel subgroup B and maximal parabolic subgroups P_a and P_b of G containing B as in [17] and [18]. In particular, we may assume that α stabilizes B , P_a and P_b .

By the remarks on p. 152 in [19], G has only two 2-blocks, the principal block B_0 and one defect-0-block (corresponding to the Steinberg character). Hence we have to verify Dade's conjecture only for the principal block B_0 .

According to a corollary of the Borel-Tits theorem [6], the normalizers of radical 2-subgroups are parabolic subgroups. The radical 2-chains of G (up to G -conjugacy) are given in Table 1.

TABLE 1. Radical 2-chains of G .

C		$N_G(C)$	$N_A(C)$
C_1	$\{1\}$	G	A
C_2	$\{1\} < O_p(P_a)$	P_a	$P_a \rtimes \langle \alpha \rangle$
C_3	$\{1\} < O_p(P_a) < O_p(B)$	B	$B \rtimes \langle \alpha \rangle$
C_4	$\{1\} < O_p(P_b)$	P_b	$P_b \rtimes \langle \alpha \rangle$
C_5	$\{1\} < O_p(P_b) < O_p(B)$	B	$B \rtimes \langle \alpha \rangle$
C_6	$\{1\} < O_p(B)$	B	$B \rtimes \langle \alpha \rangle$

Since C_5 and C_6 have the same normalizers $N_G(C_5) = N_G(C_6)$ and $N_A(C_5) = N_A(C_6)$, it follows that

$$k(N_G(C_5), B_0, d, u) = k(N_G(C_6), B_0, d, u)$$

for all $d \in \mathbb{N}$ and $u \mid 2n + 1$. Thus the contribution of C_5 and C_6 in the alternating sum of Dade's invariant conjecture is zero. So, Dade's invariant conjecture for G is equivalent with

$$(1) \quad k(G, B_0, d, u) + k(B, B_0, d, u) = k(P_a, B_0, d, u) + k(P_b, B_0, d, u)$$

for all $d \in \mathbb{N}$ and $u \mid 2n + 1$.

Theorem 5.1. *Let $\tilde{B}$ be a 2-block of $G = {}^2F_4(2^{2n+1})$, $n > 0$, with a positive defect. Then $\tilde{B}$ satisfies Dade's invariant conjecture.*

Proof: By the proceeding remarks, we can assume $\tilde{B} = B_0$. Suppose $u \mid 2n+1$ and set $t := \frac{2n+1}{u}$ and $H := \langle \alpha^t \rangle$. Let $S \in \{G, B, P_a, P_b\}$. By the character tables in [17], [18] and CHEVIE, we have $k(S, B_0, d, u) = 0$ when $d \notin \{11n+6, 14n+7, 14n+8, 15n+8, 16n+8, 17n+9, 18n+9, 20n+10, 20n+11, 20n+12, 21n+11, 22n+11, 23n+12, 24n+12\}$.

(a) If $d = 11n+6$, then we have $k(G, B_0, d, u) = \sum_{j \in \{19, 20\}} |C_{G I_j}(H)| = 2$ and $k(P_b, B_0, d, u) = \sum_{j \in \{46, 53\}} |C_{P_b I_j}(H)| = 2$ as well as $k(B, B_0, d, u) = k(P_a, B_0, d, u) = 0$ by Tables A.2 and A.16. Thus (1) holds in this case.

(b) If $d = 14n+7$, then we have

$$k(G, B_0, d, u) = |C_{G I_{18}}(H)| = 1 \text{ and } k(P_a, B_0, d, u) = |C_{P_a I_{31}}(H)| = 1$$

by Tables A.3 and A.16. Thus (1) holds in this case.

(c) If $d = 14n+8$, then we have $k(G, B_0, d, u) = \sum_{j \in \{55, 56, 57, 58\}} |C_{B I_j}(H)| = 4$ and $k(P_b, B_0, d, u) = \sum_{j \in \{44, 45, 51, 52\}} |C_{P_b I_j}(H)| = 4$ by Tables A.4 and A.16. Thus (1) holds in this case.

(d) If $d = 15n+8$, then we have $k(B, B_0, d, u) = \sum_{j \in \{51, 52, 53, 54\}} |C_{B I_j}(H)| = 2 \cdot 2^t$ and $k(P_b, B_0, d, u) = \sum_{j \in \{43, 47, 48, 49, 50, 54, 55, 56\}} |C_{P_b I_j}(H)| = 2 \cdot 2^t$ by Tables A.5 and A.16. Thus (1) holds in this case.

(e) If $d = 16n+8$, then Table A.6 and Proposition 4.2 imply, that (1) is equivalent to

$$\sum_{j \in \{42, 43\}} |C_{B I_j}(H)| = \sum_{j \in \{30, 32, 33\}} |C_{P_a I_j}(H)|.$$

By Table A.16, the sums on both sides of the above equation are equal. Thus (1) also holds in this case.

(f) If $d = 17n+9$, then Table A.7 and Proposition 4.2 imply, that (1) is equivalent to

$$\sum_{j \in J_B} |C_{B I_j}(H)| = \sum_{j \in J_{P_a}} |C_{P_a I_j}(H)|$$

with $J_B := \{44, 45\}$ and $J_{P_a} := \{34, 35\}$. By Table A.16, the sums on both sides of the above equation are equal. Thus (1) also holds in this case.

(g) If $d = 18n+9$, then Table A.8 and Proposition 4.2 imply, that (1) is equivalent to

$$|C_{G I_{31}}(H)| + |C_{B I_{39}}(H)| = |C_{P_a I_{29}}(H)| + |C_{P_b I_{40}}(H)|.$$

By Table A.16, the sums on both sides of the above equation are equal. Thus (1) also holds in this case.

(h) If $d = 20n+10$, then Table A.9 and Proposition 4.2 imply, that (1) is equivalent to

$$\sum_{j \in J_G} |C_{G I_j}(H)| + \sum_{j \in J_B} |C_{B I_j}(H)| = \sum_{j \in J_{P_a}} |C_{P_a I_j}(H)| + \sum_{j \in J_{P_b}} |C_{P_b I_j}(H)|$$

where the sums are over $J_G = \{17, 25, 45, 49\}$, $J_B = \{11, 12, 23\}$, $J_{P_a} = \{12, 20, 26\}$, $J_{P_b} = \{4, 16, 21, 26\}$, respectively. By Table A.16, we have

$$k(G, B_0, d, u) + k(B, B_0, d, u) = \sum_{j \in J_G} |C_{GI_j}(H)| + \sum_{j \in J_B} |C_{BI_j}(H)| = 4 \cdot 2^t - 3$$

and

$$k(P_a, B_0, d, u) + k(P_b, B_0, d, u) = \sum_{j \in J_{P_a}} |C_{P_a I_j}(H)| + \sum_{j \in J_{P_b}} |C_{P_b I_j}(H)| = 4 \cdot 2^t - 3.$$

Thus (1) also holds in this case.

(i) If $d = 20n + 11$, then Table A.10 and Proposition 4.2 imply, that (1) is equivalent to

$$\sum_{j \in J_G} |C_{GI_j}(H)| + \sum_{j \in J_B} |C_{BI_j}(H)| = \sum_{j \in J_{P_a}} |C_{P_a I_j}(H)| + \sum_{j \in J_{P_b}} |C_{P_b I_j}(H)|$$

where the sums are over $J_G := \{7, 10\}$, $J_B := \{13, 14\}$, $J_{P_a} := \{13, 25\}$ and $J_{P_b} := \{14, 15\}$, respectively. By Table A.16, we have

$$k(G, B_0, d, u) + k(B, B_0, d, u) = \sum_{j \in J_G} |C_{GI_j}(H)| + \sum_{j \in J_B} |C_{BI_j}(H)| = 4$$

and

$$k(P_a, B_0, d, u) + k(P_b, B_0, d, u) = \sum_{j \in J_{P_a}} |C_{P_a I_j}(H)| + \sum_{j \in J_{P_b}} |C_{P_b I_j}(H)| = 4.$$

Thus (1) also holds in this case.

(j) If $d = 20n + 12$, then Table A.11 implies, that (1) is equivalent to

$$\sum_{j \in J_G} |C_{GI_j}(H)| + \sum_{j \in J_B} |C_{BI_j}(H)| = \sum_{j \in J_{P_a}} |C_{P_a I_j}(H)| + \sum_{j \in J_{P_b}} |C_{P_b I_j}(H)|$$

with $J_G := \{5, 6, 8, 9, 11, 12, 13, 14\}$, $J_B := \{15, 16, 17, 18, 19, 20, 21, 22\}$, $J_{P_a} := \{14, 15, 16, 17, 21, 22, 23, 24\}$, $J_{P_b} := \{17, 18, 19, 20, 22, 23, 24, 25\}$. By Table A.16, we have

$$k(G, B_0, d, u) + k(B, B_0, d, u) = \sum_{j \in J_G} |C_{GI_j}(H)| + \sum_{j \in J_B} |C_{BI_j}(H)| = 16$$

and

$$k(P_a, B_0, d, u) + k(P_b, B_0, d, u) = \sum_{j \in J_{P_a}} |C_{P_a I_j}(H)| + \sum_{j \in J_{P_b}} |C_{P_b I_j}(H)| = 16.$$

Thus (1) also holds in this case.

(k) If $d = 21n + 11$, then Table A.12 implies, that (1) is equivalent to

$$\sum_{j \in J_B} |C_{BI_j}(H)| = \sum_{j \in J_{P_b}} |C_{P_b I_j}(H)|$$

with $J_B := \{26, 27, 28, 29\}$ and $J_{P_b} := \{27, 28, 29, 30\}$. By Table A.16, the sums on both sides of the above equation are equal. Thus (1) also holds in this case.

(l) If $d = 22n + 11$, then Table A.13 implies, that (1) is equivalent to

$$\sum_{j \in J_G} |C_{GI_j}(H)| + \sum_{j \in J_B} |C_{BI_j}(H)| = \sum_{j \in J_{P_a}} |C_{P_a I_j}(H)| + \sum_{j \in J_{P_b}} |C_{P_b I_j}(H)|$$

with $J_G := \{4, 27, 30, 33\}$, $J_B := \{9, 10\}$, $J_{P_a} := \{2, 11\}$ and $J_{P_b} := \{12, 13\}$. By Table A.16, we have

$$k(G, B_0, d, u) + k(B, B_0, d, u) = \sum_{j \in J_G} |C_{GI_j}(H)| + \sum_{j \in J_B} |C_{BI_j}(H)| = 2 \cdot 2^t$$

and

$$k(P_a, B_0, d, u) + k(P_b, B_0, d, u) = \sum_{j \in J_{P_a}} |C_{P_a I_j}(H)| + \sum_{j \in J_{P_b}} |C_{P_b I_j}(H)| = 2 \cdot 2^t.$$

Thus (1) also holds in this case.

(m) If $d = 23n + 12$, then Table A.14 implies, that (1) is equivalent to

$$\sum_{j \in J_G} |C_{GI_j}(H)| + \sum_{j \in J_B} |C_{BI_j}(H)| = \sum_{j \in J_{P_a}} |C_{P_a I_j}(H)| + \sum_{j \in J_{P_b}} |C_{P_b I_j}(H)|$$

with $J_G := \{2, 3, 23, 24, 43, 44, 47, 48\}$, $J_B := \{3, 4, 6, 7\}$, $J_{P_a} := \{7, 8, 9, 10\}$ and $J_{P_b} := \{2, 3, 9, 10\}$. By Table A.16, we have

$$k(G, B_0, d, u) + k(B, B_0, d, u) = \sum_{j \in J_G} |C_{GI_j}(H)| + \sum_{j \in J_B} |C_{BI_j}(H)| = 4 \cdot 2^t$$

and

$$k(P_a, B_0, d, u) + k(P_b, B_0, d, u) = \sum_{j \in J_{P_a}} |C_{P_a I_j}(H)| + \sum_{j \in J_{P_b}} |C_{P_b I_j}(H)| = 4 \cdot 2^t.$$

Thus (1) also holds in this case.

(n) If $d = 24n + 12$, then Table A.15 implies, that (1) is equivalent to

$$|C_{GI^{ss}}(H)| + \sum_{j \in J_B} |C_{BI_j}(H)| = \sum_{j \in J_{P_a}} |C_{P_a I_j}(H)| + \sum_{j \in J_{P_b}} |C_{P_b I_j}(H)|$$

with $J_B := \{1, 2, 5, 8\}$, $J_{P_a} := \{1, 3, 4, 5, 6\}$ and $J_{P_b} := \{1, 5, 6, 7, 8, 11\}$. By [17, Lemma 6.2] and Table A.16, we have

$$k(G, B_0, d, u) + k(B, B_0, d, u) = |C_{GI^{ss}}(H)| + \sum_{j \in J_B} |C_{BI_j}(H)| = 2 \cdot 2^{2t}$$

and

$$k(P_a, B_0, d, u) + k(P_b, B_0, d, u) = \sum_{j \in J_{P_a}} |C_{P_a I_j}(H)| + \sum_{j \in J_{P_b}} |C_{P_b I_j}(H)| = 2 \cdot 2^{2t}.$$

Thus (1) also holds in this case. $\square$

APPENDIX A

Table A.1 *Parameter sets for the irreducible characters of G , B , P_a , P_b . (For the parameter sets GI_{34} , GI_{35} , GI_{36} , $\dots$, GI_{41} , GI_{42} , GI_{46} , GI_{50} , GI_{51} see the remarks in Section 4. In this table, we use the abbreviations $k' := \frac{q}{\sqrt{2}}(k+l)$, $l' := \frac{q}{\sqrt{2}}(k-l)$, $a' := -\sqrt{2}q^3 + \sqrt{2}q + 1$, $b' := -\sqrt{2}q^3 - q^2 + \sqrt{2}q + 2$, $a'' := \sqrt{2}q^3 - \sqrt{2}q + 1$ and $b'' := \sqrt{2}q^3 - q^2 - \sqrt{2}q + 2$.)*

Parameter Set	Characters	Parameters	Equivalence Relation	Number of Characters
$GI_1 = \dots = GI_{21}$	$\chi_1, \dots, \chi_{21}$			1
$GI_{22} = \dots = GI_{27}$	$\chi_{22}(k), \dots, \chi_{27}(k)$	$k = 0, \dots, q^2 - 2$ $k \neq 0$	$\{k \equiv -k\}$	$\frac{q^2-2}{2}$
GI_{28}	$\chi_{28}(k, l)$	see the remarks in Section 4		$\frac{q^4-10q^2+16}{16}$
$GI_{29} = \dots = GI_{31}$	$\chi_{29}, \chi_{30}, \chi_{31}$			1
$GI_{32} = GI_{33}$	$\chi_{32}(k), \chi_{33}(k)$	$k = 0, \dots, q^2$ $\frac{q^2+1}{3} \nmid k$	$\{k \equiv -k\}$	$\frac{q^2-2}{2}$
GI_{34}	$\chi_{34}(k)$	see the remarks in Section 4		$\frac{q^4-2q^2}{4}$
GI_{35}	$\chi_{35}(k)$	see the remarks in Section 4		$\frac{1}{8}(q^4 - \sqrt{2}q^3 - 2q^2 + 2\sqrt{2}q)$
GI_{36}	$\chi_{36}(k)$	see the remarks in Section 4		$\frac{1}{8}(q^4 + \sqrt{2}q^3 - 2q^2 - 2\sqrt{2}q)$
GI_{37}	$\chi_{37}(k, l)$	see the remarks in Section 4		$\frac{q^4-2q^2}{16}$
GI_{38}	$\chi_{38}(k, l)$	see the remarks in Section 4		$\frac{q^4-10q^2+16}{48}$
GI_{39}	$\chi_{39}(k)$	see the remarks in Section 4		$\frac{q^4-q^2-2}{6}$
GI_{40}	$\chi_{40}(k)$	see the remarks in Section 4		$\frac{1}{12}(q^4 - \sqrt{2}q^3 + q^2 - \sqrt{2}q)$
GI_{41}	$\chi_{41}(k)$	see the remarks in Section 4		$\frac{1}{12}(q^4 + \sqrt{2}q^3 + q^2 + \sqrt{2}q)$
GI_{42}	$\chi_{42}(k)$	see the remarks in Section 4		$\frac{q^2-\sqrt{2}q}{4}$
$GI_{43} = \dots = GI_{45}$	$\chi_{43}(k), \dots, \chi_{45}(k)$	$k = 0, \dots, q^2 - \sqrt{2}q$ $k \neq 0$	$\{k \equiv -k \equiv q^2k \equiv -q^2k\}$	$\frac{q^2-\sqrt{2}q}{4}$
GI_{46}	$\chi_{46}(k)$	see the remarks in Section 4		$\frac{q^2+\sqrt{2}q}{4}$
$GI_{47} = \dots = GI_{49}$	$\chi_{47}(k), \dots, \chi_{49}(k)$	$k = 0, \dots, q^2 + \sqrt{2}q$ $k \neq 0$	$\{k \equiv -k \equiv q^2k \equiv -q^2k\}$	$\frac{q^2+\sqrt{2}q}{4}$
GI_{50}	$\chi_{50}(k, l)$	see the remarks in Section 4		
GI_{51}	$\chi_{51}(k, l)$	see the remarks in Section 4		
BI_1	$B\chi_1(k, l)$	$k, l = 0, \dots, q^2 - 2$		$(q^2 - 1)^2$
$BI_2 = \dots = BI_5$	$B\chi_2(k), \dots, B\chi_5(k)$	$k = 0, \dots, q^2 - 2$		$q^2 - 1$
$BI_6 = \dots = BI_8$	$B\chi_6, B\chi_7, B\chi_8$			1
BI_9	$B\chi_9(k)$	$k = 0, \dots, q^2 - 2$		$q^2 - 1$
BI_{10}	$B\chi_{10}$			1

Table A.1 (cont.) *Parameter sets for irreducible characters.*

Parameter Set	Characters	Parameters	Equivalence Relation	Number of Characters
$BI_{11} = BI_{12}$	$B\chi_{11}(k), B\chi_{12}(k)$	$k = 0, \dots, q^2 - 2$		$q^2 - 1$
$BI_{13} = \dots = BI_{22}$	$B\chi_{13}, \dots, B\chi_{22}$			1
BI_{23}	$B\chi_{23}(k)$	$k = 0, \dots, q^2 - 2$		$q^2 - 1$
BI_{24}	$B\chi_{24}$			1
BI_{25}	$B\chi_{25}(k)$	$k = 1, \dots, q^2$		q^2
$BI_{26} = BI_{27}$	$B\chi_{26}(k), B\chi_{27}(k)$	$k = 0, \dots, q^2 - 2$		$q^2 - 1$
$BI_{28} = \dots = BI_{34}$	$B\chi_{28}, \dots, B\chi_{34}$			1
BI_{35}	$B\chi_{35}(k)$	$k = 1, \dots, 4 \cdot \frac{q^2-2}{6}$		$4 \cdot \frac{q^2-2}{6}$
BI_{36}	$B\chi_{36}(k)$	$k = 1, \dots, 4 \cdot \frac{q^2-2}{2}$		$4 \cdot \frac{q^2-2}{2}$
BI_{37}	$B\chi_{37}(k)$	$k = 1, \dots, \frac{q^2+1}{3}$		$\frac{q^2+1}{3}$
BI_{38}	$B\chi_{38}$			1
BI_{39}	$B\chi_{39}(k)$	$k = 0, \dots, q^2 - 2$		$q^2 - 1$
BI_{40}	$B\chi_{40}$			1
BI_{41}	$B\chi_{41}(k)$	$k = 1, \dots, q^2$		q^2
BI_{42}	$B\chi_{42}(k)$	$k = 0, \dots, q^2 - 2$		$q^2 - 1$
BI_{43}	$B\chi_{43}$			1
$BI_{44} = BI_{45}$	$B\chi_{44}(k), B\chi_{45}(k)$	$k = 0, \dots, q^2 - 2$		$q^2 - 1$
$BI_{46} = BI_{47}$	$B\chi_{46}, B\chi_{47}$			1
$BI_{48} = \dots = BI_{50}$	$B\chi_{48}(k), \dots, B\chi_{50}(k)$	$k = 1, \dots, q^2$		q^2
$BI_{51} = BI_{52}$	$B\chi_{51}(k), B\chi_{52}(k)$	$k = 0, \dots, q^2 - 2$		$q^2 - 1$
$BI_{53} = \dots = BI_{58}$	$B\chi_{53}, \dots, B\chi_{58}$			1
$PaI_1 = PaI_2$	$Pa\chi_1(k), Pa\chi_2(k)$	$k = 0, \dots, q^2 - 2$		$q^2 - 1$
PaI_3	$Pa\chi_3(k, l)$	$k, l = 0, \dots, q^2 - 2$ $l \neq (\sqrt{2}q - 1)k$	$\{(k, l) \equiv (k', l')\}$	$\frac{q^4 - 3q^2 + 2}{2}$
PaI_4	$Pa\chi_4(k)$	$k = 0, \dots, q^4 - 2$ $q^2 + 1 \nmid k$	$\{k \equiv q^2 k\}$	$\frac{q^4 - q^2}{2}$
PaI_5	$Pa\chi_5(k)$	$k = 0, \dots, q^2 - 2$		$q^2 - 1$
PaI_6	$Pa\chi_6$			1
$PaI_7 = PaI_8$	$Pa\chi_7(k), Pa\chi_8(k)$	$k = 0, \dots, q^2 - 2$		$q^2 - 1$
$PaI_9 = \dots = PaI_{11}$	$Pa\chi_9, \dots, Pa\chi_{11}$			1
PaI_{12}	$Pa\chi_{12}(k)$	$k = 0, \dots, q^2 - 2$		$q^2 - 1$
$PaI_{13} = \dots = PaI_{25}$	$Pa\chi_{13}, \dots, Pa\chi_{25}$			1
PaI_{26}	$Pa\chi_{26}(k)$	$k = 0, \dots, q^2 - 2$		$q^2 - 1$
PaI_{27}	$Pa\chi_{27}$			1
PaI_{28}	$Pa\chi_{28}(k)$	$k = 1, \dots, q^2$		q^2
$PaI_{29} = \dots = PaI_{31}$	$Pa\chi_{29}, \dots, Pa\chi_{31}$			1
PaI_{32}	$Pa\chi_{32}(k)$	$k = 0, \dots, q^2 - 2; k \neq 0$	$\{k \equiv -k\}$	$\frac{q^2-2}{2}$
PaI_{33}	$Pa\chi_{33}(k)$	$k = 0, \dots, q^2; k \neq 0$	$\{k \equiv -k\}$	$\frac{q^2}{2}$

Table A.1 (cont.) *Parameter sets for irreducible characters.*

Parameter Set	Characters	Parameters	Equivalence Relation	Number of Characters
$P_a I_{34} = P_a I_{35}$	$P_a \chi_{34}(k), P_a \chi_{35}(k)$	$k = 0, \dots, q^2 - 2$		$q^2 - 1$
$P_a I_{36} = P_a I_{37}$	$P_a \chi_{36}, P_a \chi_{37}$			1
$P_a I_{38} = P_a I_{39}$ $= P_a I_{40}$	$P_a \chi_{38}(k), P_a \chi_{39}(k),$ $P_a \chi_{40}(k)$	$k = 1, \dots, q^2$		q^2
$P_b I_1 = \dots = P_b I_4$	$P_b \chi_1(k), \dots, P_b \chi_4(k)$	$k = 0, \dots, q^2 - 2$		$q^2 - 1$
$P_b I_5$	$P_b \chi_5(k, l)$	$k, l = 0, \dots, q^2 - 2$ $l \neq 0$	$\{(k, l) \equiv (k, -l)\}$	$\frac{q^4 - 3q^2 + 2}{2}$
$P_b I_6$	$P_b \chi_6(k)$	$k = 0, \dots, \phi_1 \phi_2 \phi'_8 - 1$ $\phi'_8 \nmid k$	$\{k \equiv q^2 k \equiv a' k \equiv b' k\}$	$\frac{1}{4}(q^4 + \sqrt{2}q^3 - q^2 - \sqrt{2}q)$
$P_b I_7$	$P_b \chi_7(k)$	$k = 0, \dots, \phi_1 \phi_2 \phi''_8 - 1$ $\phi''_8 \nmid k$	$\{k \equiv q^2 k \equiv a'' k \equiv b'' k\}$	$\frac{1}{4}(q^4 - \sqrt{2}q^3 - q^2 + \sqrt{2}q)$
$P_b I_8$	$P_b \chi_8(k)$	$k = 0, \dots, q^2 - 2$		$q^2 - 1$
$P_b I_9 = P_b I_{10}$ $= P_b I_{11}$	$P_b \chi_9, P_b \chi_{10},$ $P_b \chi_{11}$			1
$P_b I_{12}$	$P_b \chi_{12}(k)$	$k = 0, \dots, q^2 - 2$		$q^2 - 1$
$P_b I_{13} = P_b I_{14}$ $= P_b I_{15}$	$P_b \chi_{13}, P_b \chi_{14},$ $P_b \chi_{15}$			1
$P_b I_{16}$	$P_b \chi_{16}(k)$	$k = 0, \dots, q^2 - 2$ $k \neq 0$	$\{k \equiv -k\}$	$\frac{q^2 - 2}{2}$
$P_b I_{17} = \dots = P_b I_{20}$	$P_b \chi_{17}, \dots, P_b \chi_{20}$			1
$P_b I_{21}$	$P_b \chi_{21}(k)$	$k = 0, \dots, q^2 - \sqrt{2}q$ $k \neq 0$	$\{k \equiv -k \equiv q^2 k \equiv -q^2 k\}$	$\frac{q^2 - \sqrt{2}q}{4}$
$P_b I_{22} = \dots = P_b I_{25}$	$P_b \chi_{22}, \dots, P_b \chi_{25}$			1
$P_b I_{26}$	$P_b \chi_{26}(k)$	$k = 0, \dots, q^2 + \sqrt{2}q$ $k \neq 0$	$\{k \equiv -k \equiv q^2 k \equiv -q^2 k\}$	$\frac{q^2 + \sqrt{2}q}{4}$
$P_b I_{27} = P_b I_{28}$	$P_b \chi_{27}(k), P_b \chi_{28}(k)$	$k = 0, \dots, q^2 - 2$		$q^2 - 1$
$P_b I_{29} = \dots = P_b I_{35}$	$P_b \chi_{29}, \dots, P_b \chi_{35}$			1
$P_b I_{36}$	$P_b \chi_{36}(k)$	$k = 1, \dots, 4 \cdot \frac{q^2 - 2}{6}$		$4 \cdot \frac{q^2 - 2}{6}$
$P_b I_{37}$	$P_b \chi_{37}(k)$	$k = 1, \dots, 4 \cdot \frac{q^2 - 2}{2}$		$4 \cdot \frac{q^2 - 2}{2}$
$P_b I_{38}$	$P_b \chi_{38}(k)$	$k = 1, \dots, \frac{q^2 + 1}{3}$		$\frac{q^2 + 1}{3}$
$P_b I_{39}$	$P_b \chi_{39}$			1
$P_b I_{40}$	$P_b \chi_{40}(k)$	$k = 0, \dots, q^2 - 2$		$q^2 - 1$
$P_b I_{41}$	$P_b \chi_{41}$			1
$P_b I_{42}$	$P_b \chi_{42}(k)$	$k = 1, \dots, q^2$		q^2
$P_b I_{43} = \dots = P_b I_{46}$	$P_b \chi_{43}, \dots, P_b \chi_{46}$			1
$P_b I_{47}$	$P_b \chi_{47}(k)$	$k = 0, \dots, q^2 - 2$ $k \neq 0$	$\{k \equiv -k\}$	$\frac{q^2 - 2}{2}$
$P_b I_{48}$	$P_b \chi_{48}(k)$	$k = 0, \dots, q^2 + \sqrt{2}q$ $k \neq 0$	$\{k \equiv -k \equiv q^2 k \equiv -q^2 k\}$	$\frac{q^2 + \sqrt{2}q}{4}$

Table A.1 (cont.) *Parameter sets for irreducible characters.*

Parameter Set	Characters	Parameters	Equivalence Relation	Number of Characters
$P_b I_{49}$	$P_b \chi_{49}(k)$	$k = 0, \dots, q^2 - \sqrt{2}q$ $k \neq 0$	$\{k \equiv -k \equiv q^2 k \equiv -q^2 k\}$	$\frac{q^2 - \sqrt{2}q}{4}$
$P_b I_{50} = \dots = P_b I_{53}$	$P_b \chi_{50}, \dots, P_b \chi_{53}$			1
$P_b I_{54}$	$P_b \chi_{54}(k)$	$k = 0, \dots, q^2 - 2$ $k \neq 0$	$\{k \equiv -k\}$	$\frac{q^2 - 2}{2}$
$P_b I_{55}$	$P_b \chi_{55}(k)$	$k = 0, \dots, q^2 + \sqrt{2}q$ $k \neq 0$	$\{k \equiv -k \equiv q^2 k \equiv -q^2 k\}$	$\frac{q^2 + \sqrt{2}q}{4}$
$P_b I_{56}$	$P_b \chi_{56}(k)$	$k = 0, \dots, q^2 - \sqrt{2}q$ $k \neq 0$	$\{k \equiv -k \equiv q^2 k \equiv -q^2 k\}$	$\frac{q^2 - \sqrt{2}q}{4}$

Table A.2 *The irreducible characters of the chain normalizers of defect $11n + 6$.*

Group	Character	Degree	Parameter	Number	Class
G	χ_{19}	$\frac{q^{13}}{\sqrt{2}} \phi_1 \phi_2 \phi_4^2 \phi_{12}$	$G I_{19}$	1	$c_{1,3}$
	χ_{20}	$\frac{q^{13}}{\sqrt{2}} \phi_1 \phi_2 \phi_4^2 \phi_{12}$	$G I_{20}$	1	$c_{1,3}$
P_b	$P_b \chi_{46}$	$\frac{q^{13}}{\sqrt{2}} \phi_1 \phi_2$	$P_b I_{46}$	1	$c_{1,9}$
	$P_b \chi_{53}$	$\frac{q^{13}}{\sqrt{2}} \phi_1 \phi_2$	$P_b I_{53}$	1	$c_{1,9}$

Table A.3 *The irreducible characters of the chain normalizers of defect $14n + 7$.*

Group	Character	Degree	Parameter	Number	Class
G	χ_{18}	$q^{10} \phi_{12} \phi_{24}$	$G I_{18}$	1	
P_a	$P_a \chi_{31}$	$q^{10} \phi_1 \phi_2$	$P_a I_{31}$	1	

Table A.4 *The irreducible characters of the chain normalizers of defect $14n + 8$.*

Group	Character	Degree	Parameter	Number	Class
B	$B \chi_{55}$	$\frac{1}{2} q^{10} \phi_1^2 \phi_2^2$	$B I_{55}$	1	$c_{1,57}, c_{1,62}$
	$B \chi_{56}$	$\frac{1}{2} q^{10} \phi_1^2 \phi_2^2$	$B I_{56}$	1	$c_{1,57}, c_{1,62}$
	$B \chi_{57}$	$\frac{1}{2} q^{10} \phi_1^2 \phi_2^2$	$B I_{57}$	1	$c_{1,57}, c_{1,62}$
	$B \chi_{58}$	$\frac{1}{2} q^{10} \phi_1^2 \phi_2^2$	$B I_{58}$	1	$c_{1,57}, c_{1,62}$
P_b	$P_b \chi_{44}$	$\frac{1}{2} q^{10} \phi_1^2 \phi_2^2$	$P_b I_{44}$	1	$c_{1,31}, c_{1,33}$
	$P_b \chi_{45}$	$\frac{1}{2} q^{10} \phi_1^2 \phi_2^2$	$P_b I_{45}$	1	$c_{1,31}, c_{1,33}$
	$P_b \chi_{51}$	$\frac{1}{2} q^{10} \phi_1^2 \phi_2^2$	$P_b I_{51}$	1	$c_{1,31}, c_{1,33}$
	$P_b \chi_{52}$	$\frac{1}{2} q^{10} \phi_1^2 \phi_2^2$	$P_b I_{52}$	1	$c_{1,31}, c_{1,33}$

Table A.5 *The irreducible characters of the chain normalizers of defect $15n + 8$.*

Group	Character	Degree	Parameter	Number	Class
B	$B\chi_{51}(k)$	$\frac{q^9}{\sqrt{2}}\phi_1\phi_2$	$B I_{51}$	$q^2 - 1$	$c_{5,0}(i), c_{5,2}(i)$
	$B\chi_{52}(k)$	$\frac{q^9}{\sqrt{2}}\phi_1\phi_2$	$B I_{52}$	$q^2 - 1$	$c_{5,0}(i), c_{5,2}(i)$
	$B\chi_{53}$	$\frac{q^9}{\sqrt{2}}\phi_1^2\phi_2^2$	$B I_{53}$	1	$c_{1,12}$
	$B\chi_{54}$	$\frac{q^9}{\sqrt{2}}\phi_1^2\phi_2^2$	$B I_{54}$	1	$c_{1,12}$
P_b	$P_b\chi_{43}$	$\frac{q^9}{\sqrt{2}}\phi_1\phi_2$	$P_b I_{43}$	1	$c_{1,9}$
	$P_b\chi_{47}(k)$	$\frac{q^9}{\sqrt{2}}\phi_1\phi_2\phi_8$	$P_b I_{47}$	$\frac{1}{2}(q^2 - 2)$	$c_{5,0}(i), c_{5,2}(i)$
	$P_b\chi_{48}(k)$	$\frac{q^9}{\sqrt{2}}\phi_1^2\phi_2^2\phi_8''$	$P_b I_{48}$	$\frac{1}{4}(q^2 + \sqrt{2}q)$	$c_{10,0}(i), c_{10,2}(i)$
	$P_b\chi_{49}(k)$	$\frac{q^9}{\sqrt{2}}\phi_1^2\phi_2^2\phi_8'$	$P_b I_{49}$	$\frac{1}{4}(q^2 - \sqrt{2}q)$	$c_{8,0}(i), c_{8,2}(i)$
	$P_b\chi_{50}$	$\frac{q^9}{\sqrt{2}}\phi_1\phi_2$	$P_b I_{50}$	1	$c_{1,9}$
	$P_b\chi_{54}(k)$	$\frac{q^9}{\sqrt{2}}\phi_1\phi_2\phi_8$	$P_b I_{54}$	$\frac{1}{2}(q^2 - 2)$	$c_{5,0}(i), c_{5,2}(i)$
	$P_b\chi_{55}(k)$	$\frac{q^9}{\sqrt{2}}\phi_1^2\phi_2^2\phi_8''$	$P_b I_{55}$	$\frac{1}{4}(q^2 + \sqrt{2}q)$	$c_{10,0}(i), c_{10,2}(i)$
	$P_b\chi_{56}(k)$	$\frac{q^9}{\sqrt{2}}\phi_1^2\phi_2^2\phi_8'$	$P_b I_{56}$	$\frac{1}{4}(q^2 - \sqrt{2}q)$	$c_{8,0}(i), c_{8,2}(i)$

Table A.6 *The irreducible characters of the chain normalizers of defect $16n + 8$.*

Group	Character	Degree	Parameter	Number	Class
B	$B\chi_{42}(k)$	$q^8\phi_1\phi_2$	$B I_{42}$	$q^2 - 1$	$c_{9,0}(i)$
	$B\chi_{43}$	$q^8\phi_1^2\phi_2^2$	$B I_{43}$	1	$c_{1,2}$
	$B\chi_{50}(k)$	$q^8\phi_1^2\phi_2^2$	$B I_{50}$	q^2	
P_a	$P_a\chi_{30}$	$q^8\phi_1\phi_2$	$P_a I_{30}$	1	
	$P_a\chi_{32}(k)$	$q^8\phi_1\phi_2\phi_4$	$P_a I_{32}$	$\frac{q^2-2}{2}$	$c_{6,0}(i)$
	$P_a\chi_{33}(k)$	$q^8\phi_1^2\phi_2^2$	$P_a I_{33}$	$\frac{q^2}{2}$	$c_{9,0}(i)$
	$P_a\chi_{40}(k)$	$q^8\phi_1^2\phi_2^2\phi_4$	$P_a I_{40}$	q^2	

Table A.7 *The irreducible characters of the chain normalizers of defect $17n + 9$.*

Group	Character	Degree	Parameter	Number	Class
B	$B\chi_{44}(k)$	$\frac{q^7}{\sqrt{2}}\phi_1\phi_2$	$B I_{44}$	$q^2 - 1$	$c_{6,0}(i), c_{6,2}(i)$
	$B\chi_{45}(k)$	$\frac{q^7}{\sqrt{2}}\phi_1\phi_2$	$B I_{45}$	$q^2 - 1$	$c_{6,0}(i), c_{6,2}(i)$
	$B\chi_{46}$	$\frac{q^7}{\sqrt{2}}\phi_1^2\phi_2^2$	$B I_{46}$	1	$c_{1,9}$
	$B\chi_{47}$	$\frac{q^7}{\sqrt{2}}\phi_1^2\phi_2^2$	$B I_{47}$	1	$c_{1,9}$
	$B\chi_{48}(k)$	$\frac{q^7}{\sqrt{2}}\phi_1^2\phi_2^2$	$B I_{48}$	q^2	
	$B\chi_{49}(k)$	$\frac{q^7}{\sqrt{2}}\phi_1^2\phi_2^2$	$B I_{49}$	q^2	
P_a	$P_a\chi_{34}(k)$	$\frac{q^7}{\sqrt{2}}\phi_1\phi_2\phi_4$	$P_a I_{34}$	$q^2 - 1$	$c_{4,0}(i), c_{4,2}(i)$
	$P_a\chi_{35}(k)$	$\frac{q^7}{\sqrt{2}}\phi_1\phi_2\phi_4$	$P_a I_{35}$	$q^2 - 1$	$c_{4,0}(i), c_{4,2}(i)$
	$P_a\chi_{36}$	$\frac{q^7}{\sqrt{2}}\phi_1^2\phi_2^2\phi_4$	$P_a I_{36}$	1	$c_{1,7}$
	$P_a\chi_{37}$	$\frac{q^7}{\sqrt{2}}\phi_1^2\phi_2^2\phi_4$	$P_a I_{37}$	1	$c_{1,7}$
	$P_a\chi_{38}(k)$	$\frac{q^7}{\sqrt{2}}\phi_1^2\phi_2^2\phi_4$	$P_a I_{38}$	q^2	
	$P_a\chi_{39}(k)$	$\frac{q^7}{\sqrt{2}}\phi_1^2\phi_2^2\phi_4$	$P_a I_{39}$	q^2	

Table A.8 *The irreducible characters of the chain normalizers of defect $18n + 9$.*

Group	Character	Degree	Parameter	Number	Class
G	χ_{31}	$q^6 \phi_1 \phi_2 \phi_8^2 \phi_{24}$	$_G I_{31}$	1	
B	$_B \chi_{38}$	$q^6 \phi_1^2 \phi_2^2$	$_B I_{38}$	1	$c_{1,4}$
	$_B \chi_{39}(k)$	$q^6 \phi_1 \phi_2$	$_B I_{39}$	$q^2 - 1$	$c_{8,0}(i)$
	$_B \chi_{40}$	$q^6 \phi_1^2 \phi_2^2$	$_B I_{40}$	1	$c_{1,4}$
	$_B \chi_{41}(k)$	$q^6 \phi_1^2 \phi_2^2$	$_B I_{41}$	q^2	
P_a	$P_a \chi_{29}$	$q^6 \phi_1^2 \phi_2^2 \phi_4$		1	
P_b	$P_b \chi_{39}$	$q^6 \phi_1^2 \phi_2^2 \phi_8$	$P_b I_{39}$	1	$c_{1,3}$
	$P_b \chi_{40}(k)$	$q^6 \phi_1 \phi_2 \phi_8$	$P_b I_{40}$	$q^2 - 1$	$c_{6,0}(i)$
	$P_b \chi_{41}$	$q^6 \phi_1^2 \phi_2^2 \phi_8$	$P_b I_{41}$	1	$c_{1,3}$
	$P_b \chi_{42}(k)$	$q^6 \phi_1^2 \phi_2^2 \phi_8$	$P_b I_{42}$	q^2	

Table A.9 *The irreducible characters of the chain normalizers of defect $20n + 10$.*

Group	Character	Degree	Parameter	Number	Class
G	χ_{15}	$\frac{1}{3} q^4 \phi_1^2 \phi_2^2 \phi_4^2 \phi_8^2$	$_G I_{15}$	1	$c_{5,3}$
	χ_{16}	$\frac{1}{3} q^4 \phi_1^2 \phi_2^2 \phi_4^2 \phi_8^2$	$_G I_{16}$	1	$c_{5,3}$
	χ_{17}	$\frac{1}{3} q^4 \phi_1^2 \phi_2^2 \phi_{12} \phi_{24}$	$_G I_{17}$	1	
	$\chi_{25}(k)$	$q^4 \phi_4^2 \phi_8 \phi_{12} \phi_{24}$	$_G I_{25}$	$\frac{1}{2}(q^2 - 2)$	$c_{2,0}(i)$
	$\chi_{45}(k)$	$q^4 \phi_1 \phi_2 \phi_4^2 \phi_8' \phi_{12} \phi_{24}$	$_G I_{45}$	$\frac{1}{4}(q^2 - \sqrt{2}q)$	$c_{8,2}(i)$
	$\chi_{49}(k)$	$q^4 \phi_1 \phi_2 \phi_4^2 \phi_8'' \phi_{12} \phi_{24}$	$_G I_{49}$	$\frac{1}{4}(q^2 + \sqrt{2}q)$	$c_{10,2}(i)$
B	$_B \chi_{11}(k)$	$q^4 \phi_1 \phi_2$	$_B I_{11}$	$q^2 - 1$	$c_{5,0}(i)$
	$_B \chi_{12}(k)$	$q^4 \phi_1 \phi_2$	$_B I_{12}$	$q^2 - 1$	$c_{6,0}(i)$
	$_B \chi_{23}(k)$	$q^4 \phi_1 \phi_2$	$_B I_{23}$	$q^2 - 1$	$c_{7,0}(i)$
	$_B \chi_{24}$	$q^4 \phi_1^2 \phi_2^2$	$_B I_{24}$	1	$c_{1,12}$
	$_B \chi_{25}(k)$	$q^4 \phi_1^2 \phi_2^2$	$_B I_{25}$	q^2	
	$_B \chi_{34}$	$q^4 \phi_1^2 \phi_2^2$	$_B I_{34}$	1	$c_{1,12}$
	$_B \chi_{37}(k)$	$q^4 \phi_1^2 \phi_2^2$	$_B I_{37}$	$\frac{1}{3}(q^2 + 1)$	
P_a	$P_a \chi_{12}(k)$	$q^4 \phi_1 \phi_2 \phi_4$	$P_a I_{12}$	$q^2 - 1$	$c_{4,0}(i)$
	$P_a \chi_{18}$	$\frac{q^4}{3} \phi_1^2 \phi_2^2 \phi_4$	$P_a I_{18}$	1	$c_{8,3}$
	$P_a \chi_{19}$	$\frac{q^4}{3} \phi_1^2 \phi_2^2 \phi_4$	$P_a I_{19}$	1	$c_{8,3}$
	$P_a \chi_{20}$	$\frac{q^4}{3} \phi_1^2 \phi_2^2 \phi_4$	$P_a I_{20}$	1	$c_{8,0}$
	$P_a \chi_{26}(k)$	$q^4 \phi_1 \phi_2 \phi_4$	$P_a I_{26}$	$q^2 - 1$	$c_{5,0}(i)$
	$P_a \chi_{27}$	$q^4 \phi_1^2 \phi_2^2 \phi_4$	$P_a I_{27}$	1	
	$P_a \chi_{28}(k)$	$q^4 \phi_1^2 \phi_2^2 \phi_4$	$P_a I_{28}$	q^2	

Table A.9 (cont.) *Irreducible characters of the chain normalizers of defect $20n + 10$.*

Group	Character	Degree	Parameter	Number	Class
P_b	$P_b \chi_4(k)$	q^4	$P_b I_4$	$q^2 - 1$	$c_{2,0}(i)$
	$P_b \chi_{16}(k)$	$q^4 \phi_1 \phi_2 \phi_8$	$P_b I_{16}$	$\frac{1}{2}(q^2 - 2)$	$c_{5,0}(i)$
	$P_b \chi_{21}(k)$	$q^4 \phi_1^2 \phi_2^2 \phi_8'$	$P_b I_{21}$	$\frac{1}{4}(q^2 - \sqrt{2}q)$	$c_{8,0}(i)$
	$P_b \chi_{26}(k)$	$q^4 \phi_1^2 \phi_2^2 \phi_8''$	$P_b I_{26}$	$\frac{1}{4}(q^2 + \sqrt{2}q)$	$c_{10,0}(i)$
	$P_b \chi_{35}$	$q^4 \phi_1^2 \phi_2^2 \phi_8$	$P_b I_{35}$	1	
	$P_b \chi_{38}(k)$	$q^4 \phi_1^2 \phi_2^2 \phi_8$	$P_b I_{38}$	$\frac{1}{3}(q^2 + 1)$	

Table A.10 *The irreducible characters of the chain normalizers of defect $20n + 11$.*

Group	Character	Degree	Parameter	Number	Class
G	χ_7	$\frac{q^4}{2} \phi_8^2 \phi_{24}$	$G I_7$	1	
	χ_{10}	$\frac{q^4}{6} \phi_1^2 \phi_2^2 \phi_4^2 \phi_{24}$	$G I_{10}$	1	
B	$B \chi_{13}$	$\frac{q^4}{2} \phi_1^2 \phi_2^2$	$B I_{13}$	1	$c_{1,5}, c_{1,38}$
	$B \chi_{14}$	$\frac{q^4}{2} \phi_1^2 \phi_2^2$	$B I_{14}$	1	$c_{1,5}, c_{1,38}$
	$B \chi_{30}$	$\frac{q^4}{2} \phi_1^2 \phi_2^2$	$B I_{30}$	1	$c_{1,17}, c_{1,54}$
	$B \chi_{31}$	$\frac{q^4}{2} \phi_1^2 \phi_2^2$	$B I_{31}$	1	$c_{1,17}, c_{1,54}$
	$B \chi_{32}$	$\frac{q^4}{2} \phi_1^2 \phi_2^2$	$B I_{32}$	1	$c_{1,17}, c_{1,54}$
	$B \chi_{33}$	$\frac{q^4}{2} \phi_1^2 \phi_2^2$	$B I_{33}$	1	$c_{1,17}, c_{1,54}$
	$B \chi_{35}(k)$	$\frac{q^4}{2} \phi_1^2 \phi_2^2$	$B I_{35}$	$\frac{1}{3}(2q^2 - 4)$	
	$B \chi_{36}(k)$	$\frac{q^4}{2} \phi_1^2 \phi_2^2$	$B I_{36}$	$2q^2 - 4$	
P_a	$P_a \chi_{13}$	$\frac{q^4}{2} \phi_1^2 \phi_2^2 \phi_4$	$P_a I_{13}$	1	
	$P_a \chi_{25}$	$\frac{q^4}{6} \phi_1^2 \phi_2^2 \phi_4$	$P_a I_{25}$	1	
P_b	$P_b \chi_{14}$	$\frac{q^4}{2} \phi_1 \phi_2 \phi_8$	$P_b I_{14}$	1	$c_{1,14}$
	$P_b \chi_{15}$	$\frac{q^4}{2} \phi_1 \phi_2 \phi_8$	$P_b I_{15}$	1	$c_{1,14}$
	$P_b \chi_{31}$	$\frac{q^4}{2} \phi_1^2 \phi_2^2 \phi_8$	$P_b I_{31}$	1	$c_{1,6}, c_{1,30}$
	$P_b \chi_{32}$	$\frac{q^4}{2} \phi_1^2 \phi_2^2 \phi_8$	$P_b I_{32}$	1	$c_{1,6}, c_{1,30}$
	$P_b \chi_{33}$	$\frac{q^4}{2} \phi_1^2 \phi_2^2 \phi_8$	$P_b I_{33}$	1	$c_{1,6}, c_{1,30}$
	$P_b \chi_{34}$	$\frac{q^4}{2} \phi_1^2 \phi_2^2 \phi_8$	$P_b I_{34}$	1	$c_{1,6}, c_{1,30}$
	$P_b \chi_{36}(k)$	$\frac{q^4}{2} \phi_1^2 \phi_2^2 \phi_8$	$P_b I_{36}$	$\frac{1}{3}(2q^2 - 4)$	
	$P_b \chi_{37}(k)$	$\frac{q^4}{2} \phi_1^2 \phi_2^2 \phi_8$	$P_b I_{37}$	$2q^2 - 4$	

Table A.11 *The irreducible characters of the chain normalizers of defect $20n + 12$.*

Group	Character	Degree	Parameter	Number	Class
G	χ_5	$\frac{q^4}{4}\phi_4^2\phi_8''^2\phi_{12}\phi_{24}'$	GI_5	1	
	χ_6	$\frac{q^4}{4}\phi_4^2\phi_8'^2\phi_{12}\phi_{24}''$	GI_6	1	
	χ_8	$\frac{q^4}{12}\phi_1^2\phi_2^2\phi_8'^2\phi_{12}\phi_{24}'$	GI_8	1	
	χ_9	$\frac{q^4}{12}\phi_1^2\phi_2^2\phi_8''^2\phi_{12}\phi_{24}''$	GI_9	1	
	χ_{11}	$\frac{q^4}{4}\phi_1^2\phi_2^2\phi_4^2\phi_{12}\phi_{24}''$	GI_{11}	1	$c_{1,3}$
	χ_{12}	$\frac{q^4}{4}\phi_1^2\phi_2^2\phi_4^2\phi_{12}\phi_{24}'$	GI_{12}	1	$c_{1,3}$
	χ_{13}	$\frac{q^4}{4}\phi_1^2\phi_2^2\phi_4^2\phi_{12}\phi_{24}'$	GI_{13}	1	$c_{1,3}$
	χ_{14}	$\frac{q^4}{4}\phi_1^2\phi_2^2\phi_4^2\phi_{12}\phi_{24}'$	GI_{14}	1	$c_{1,3}$
B	$B\chi_{15}$	$\frac{q^4}{4}\phi_1^2\phi_2^2$	BI_{15}	1	$c_{1,23}, c_{1,38}, c_{1,54}$
	$B\chi_{16}$	$\frac{q^4}{4}\phi_1^2\phi_2^2$	BI_{16}	1	$c_{1,23}, c_{1,38}, c_{1,54}$
	$B\chi_{17}$	$\frac{q^4}{4}\phi_1^2\phi_2^2$	BI_{17}	1	$c_{1,23}, c_{1,38}, c_{1,54}$
	$B\chi_{18}$	$\frac{q^4}{4}\phi_1^2\phi_2^2$	BI_{18}	1	$c_{1,23}, c_{1,38}, c_{1,54}$
	$B\chi_{19}$	$\frac{q^4}{4}\phi_1^2\phi_2^2$	BI_{19}	1	$c_{1,23}, c_{1,38}, c_{1,54}$
	$B\chi_{20}$	$\frac{q^4}{4}\phi_1^2\phi_2^2$	BI_{20}	1	$c_{1,23}, c_{1,38}, c_{1,54}$
	$B\chi_{21}$	$\frac{q^4}{4}\phi_1^2\phi_2^2$	BI_{21}	1	$c_{1,23}, c_{1,38}, c_{1,54}$
	$B\chi_{22}$	$\frac{q^4}{4}\phi_1^2\phi_2^2$	BI_{22}	1	$c_{1,23}, c_{1,38}, c_{1,54}$
P_a	$P_a\chi_{14}$	$\frac{q^4}{4}\phi_1^2\phi_2^2\phi_4$	P_aI_{14}	1	$c_{1,37}, c_{1,39}$
	$P_a\chi_{15}$	$\frac{q^4}{4}\phi_1^2\phi_2^2\phi_4$	P_aI_{15}	1	$c_{1,37}, c_{1,39}$
	$P_a\chi_{16}$	$\frac{q^4}{4}\phi_1^2\phi_2^2\phi_4$	P_aI_{16}	1	$c_{1,37}, c_{1,39}$
	$P_a\chi_{17}$	$\frac{q^4}{4}\phi_1^2\phi_2^2\phi_4$	P_aI_{17}	1	$c_{1,37}, c_{1,39}$
	$P_a\chi_{21}$	$\frac{q^4}{4}\phi_1^2\phi_2^2\phi_4$	P_aI_{21}	1	$c_{1,39}$
	$P_a\chi_{22}$	$\frac{q^4}{4}\phi_1^2\phi_2^2\phi_4$	P_aI_{22}	1	$c_{1,39}$
	$P_a\chi_{23}$	$\frac{q^4}{12}\phi_1^2\phi_2^2\phi_4$	P_aI_{23}	1	$c_{1,18}$
	$P_a\chi_{24}$	$\frac{q^4}{12}\phi_1^2\phi_2^2\phi_4$	P_aI_{24}	1	$c_{1,18}$
P_b	$P_b\chi_{17}$	$\frac{q^4}{4}\phi_1^2\phi_2^2\phi_8'$	P_bI_{17}	1	$c_{1,14}, c_{1,30}$
	$P_b\chi_{18}$	$\frac{q^4}{4}\phi_1^2\phi_2^2\phi_8'$	P_bI_{18}	1	$c_{1,14}, c_{1,30}$
	$P_b\chi_{19}$	$\frac{q^4}{4}\phi_1^2\phi_2^2\phi_8'$	P_bI_{19}	1	$c_{1,14}, c_{1,30}$
	$P_b\chi_{20}$	$\frac{q^4}{4}\phi_1^2\phi_2^2\phi_8'$	P_bI_{20}	1	$c_{1,14}, c_{1,30}$
	$P_b\chi_{22}$	$\frac{q^4}{4}\phi_1^2\phi_2^2\phi_8''$	P_bI_{22}	1	$c_{1,14}, c_{1,30}$
	$P_b\chi_{23}$	$\frac{q^4}{4}\phi_1^2\phi_2^2\phi_8''$	P_bI_{23}	1	$c_{1,14}, c_{1,30}$
	$P_b\chi_{24}$	$\frac{q^4}{4}\phi_1^2\phi_2^2\phi_8''$	P_bI_{24}	1	$c_{1,14}, c_{1,30}$
	$P_b\chi_{25}$	$\frac{q^4}{4}\phi_1^2\phi_2^2\phi_8''$	P_bI_{25}	1	$c_{1,14}, c_{1,30}$

Table A.12 *The irreducible characters of the chain normalizers of defect $21n + 11$.*

Group	Character	Degree	Parameter	Number	Class
B	$B\chi_{26}(k)$	$\frac{q^3}{\sqrt{2}}\phi_1\phi_2$	$B I_{26}$	$q^2 - 1$	$c_{4,0}(i), c_{4,2}(i)$
	$B\chi_{27}(k)$	$\frac{q^3}{\sqrt{2}}\phi_1\phi_2$	$B I_{27}$	$q^2 - 1$	$c_{4,0}(i), c_{4,2}(i)$
	$B\chi_{28}$	$\frac{q^3}{\sqrt{2}}\phi_1^2\phi_2^2$	$B I_{28}$	1	$c_{1,17}$
	$B\chi_{29}$	$\frac{q^3}{\sqrt{2}}\phi_1^2\phi_2^2$	$B I_{29}$	1	$c_{1,17}$
P_b	$P_b\chi_{27}(k)$	$\frac{q^3}{\sqrt{2}}\phi_1\phi_2\phi_8$	$P_b I_{27}$	$q^2 - 1$	$c_{4,0}(i), c_{4,2}(i)$
	$P_b\chi_{28}(k)$	$\frac{q^3}{\sqrt{2}}\phi_1\phi_2\phi_8$	$P_b I_{28}$	$q^2 - 1$	$c_{4,0}(i), c_{4,2}(i)$
	$P_b\chi_{29}$	$\frac{q^3}{\sqrt{2}}\phi_1^2\phi_2^2\phi_8$	$P_b I_{29}$	1	$c_{1,6}$
	$P_b\chi_{30}$	$\frac{q^3}{\sqrt{2}}\phi_1^2\phi_2^2\phi_8$	$P_b I_{30}$	1	$c_{1,6}$

Table A.13 *The irreducible characters of the chain normalizers of defect $22n + 11$.*

Group	Character	Degree	Parameter	Number	Class
G	χ_4	$q^2\phi_{12}\phi_{24}$	$G I_4$	1	$c_{2,0}(i)$
	$\chi_{27}(k)$	$q^2\phi_4\phi_8^2\phi_{12}\phi_{24}$	$G I_{27}$	$\frac{1}{2}(q^2 - 2)$	
	χ_{30}	$q^2\phi_1^2\phi_2^2\phi_8^2\phi_{24}$	$G I_{30}$	1	$c_{6,0}(i)$
	$\chi_{33}(k)$	$q^2\phi_1\phi_2\phi_8^2\phi_{12}\phi_{24}$	$G I_{33}$	$\frac{1}{2}(q^2 - 2)$	
B	$B\chi_9(k)$	$q^2\phi_1\phi_2$	$B I_9$	$q^2 - 1$	$c_{4,0}(i)$
	$B\chi_{10}$	$q^2\phi_1^2\phi_2^2$	$B I_{10}$	1	
P_a	$P_a\chi_2(k)$	q^2	$P_a I_2$	$q^2 - 1$	$c_{2,0}(i)$
	$P_a\chi_{11}$	$q^2\phi_1^2\phi_2^2\phi_4$	$P_a I_{11}$	1	
P_b	$P_b\chi_{12}(k)$	$q^2\phi_1\phi_2\phi_8$	$P_b I_{12}$	$q^2 - 1$	$c_{4,0}(i)$
	$P_b\chi_{13}$	$q^2\phi_1^2\phi_2^2\phi_8$	$P_b I_{13}$	1	

Table A.14 *The irreducible characters of the chain normalizers of defect $23n + 12$.*

Group	Character	Degree	Parameter	Number	Class
G	χ_2	$\frac{q}{\sqrt{2}}\phi_1\phi_2\phi_4^2\phi_{12}$	$G I_2$	1	$c_{1,4}$
	χ_3	$\frac{q}{\sqrt{2}}\phi_1\phi_2\phi_4^2\phi_{12}$	$G I_3$	1	$c_{1,4}$
	$\chi_{23}(k)$	$\frac{q}{\sqrt{2}}\phi_1\phi_2\phi_4^2\phi_8\phi_{12}\phi_{24}$	$G I_{23}$	$\frac{1}{2}(q^2 - 2)$	$c_{2,0}(i), c_{2,2}(i)$
	$\chi_{24}(k)$	$\frac{q}{\sqrt{2}}\phi_1\phi_2\phi_4^2\phi_8\phi_{12}\phi_{24}$	$G I_{24}$	$\frac{1}{2}(q^2 - 2)$	$c_{2,0}(i), c_{2,2}(i)$
	$\chi_{43}(k)$	$\frac{q}{\sqrt{2}}\phi_1^2\phi_2^2\phi_4^2\phi_8'\phi_{12}\phi_{24}$	$G I_{43}$	$\frac{1}{4}(q^2 - \sqrt{2}q)$	$c_{1,11}, c_{8,2}(i)$
	$\chi_{44}(k)$	$\frac{q}{\sqrt{2}}\phi_1^2\phi_2^2\phi_4^2\phi_8'\phi_{12}\phi_{24}$	$G I_{44}$	$\frac{1}{4}(q^2 - \sqrt{2}q)$	$c_{1,11}, c_{8,2}(i)$
	$\chi_{47}(k)$	$\frac{q}{\sqrt{2}}\phi_1^2\phi_2^2\phi_4^2\phi_8''\phi_{12}\phi_{24}$	$G I_{47}$	$\frac{1}{4}(q^2 + \sqrt{2}q)$	$c_{1,11}, c_{10,2}(i)$
	$\chi_{48}(k)$	$\frac{q}{\sqrt{2}}\phi_1^2\phi_2^2\phi_4^2\phi_8''\phi_{12}\phi_{24}$	$G I_{48}$	$\frac{1}{4}(q^2 + \sqrt{2}q)$	$c_{1,11}, c_{10,2}(i)$
B	$B\chi_3(k)$	$\frac{q}{\sqrt{2}}\phi_1\phi_2$	$B I_3$	$q^2 - 1$	$c_{2,0}(i), c_{2,2}(i)$
	$B\chi_4(k)$	$\frac{q}{\sqrt{2}}\phi_1\phi_2$	$B I_4$	$q^2 - 1$	$c_{2,0}(i), c_{2,2}(i)$
	$B\chi_6$	$\frac{q}{\sqrt{2}}\phi_1^2\phi_2^2$	$B I_6$	1	$c_{1,54}$
	$B\chi_7$	$\frac{q}{\sqrt{2}}\phi_1^2\phi_2^2$	$B I_7$	1	$c_{1,54}$

Table A.14 (cont.) Irreducible characters of the chain normalizers of defect $23n+12$.

Group	Character	Degree	Parameter	Number	Class
P_a	$P_a \chi_7(k)$	$\frac{q}{\sqrt{2}}\phi_1\phi_2\phi_4$	$P_a I_7$	$q^2 - 1$	$c_{3,0}(i), c_{3,2}(i)$
	$P_a \chi_8(k)$	$\frac{q}{\sqrt{2}}\phi_1\phi_2\phi_4$	$P_a I_8$	$q^2 - 1$	$c_{3,0}(i), c_{3,2}(i)$
	$P_a \chi_9$	$\frac{q}{\sqrt{2}}\phi_1^2\phi_2^2\phi_4$	$P_a I_9$	1	$c_{1,12}$
	$P_a \chi_{10}$	$\frac{q}{\sqrt{2}}\phi_1^2\phi_2^2\phi_4$	$P_a I_{10}$	1	$c_{1,12}$
P_b	$P_b \chi_2(k)$	$\frac{q}{\sqrt{2}}\phi_1\phi_2$	$P_b I_2$	$q^2 - 1$	$c_{2,0}(i), c_{2,2}(i)$
	$P_b \chi_3(k)$	$\frac{q}{\sqrt{2}}\phi_1\phi_2$	$P_b I_3$	$q^2 - 1$	$c_{2,0}(i), c_{2,2}(i)$
	$P_b \chi_9$	$\frac{q}{\sqrt{2}}\phi_1^2\phi_2^2\phi_8$	$P_b I_9$	1	$c_{1,30}$
	$P_b \chi_{10}$	$\frac{q}{\sqrt{2}}\phi_1^2\phi_2^2\phi_8$	$P_b I_{10}$	1	$c_{1,30}$

Table A.15 The irreducible characters of the chain normalizers of defect $24n + 12$.
(The parameter set $G I^{ss}$ parameterizes the semisimple irreducible characters of G .)

Group	Character	Degree	Parameter	Number	Class
G			$G I^{ss}$	q^4	
B	$B \chi_1(k, l)$	1	$B I_1$	$(q^2 - 1)^2$	$c_{10,0}(i, j)$
	$B \chi_2(k)$	$\phi_1\phi_2$	$B I_2$	$q^2 - 1$	$c_{2,0}(i)$
	$B \chi_5(k)$	$\phi_1\phi_2$	$B I_5$	$q^2 - 1$	$c_{3,0}(i)$
	$B \chi_8$	$\phi_1^2\phi_2^2$	$B I_8$	1	
P_a	$P_a \chi_1(k)$	1	$P_a I_1$	$q^2 - 1$	$c_{2,0}(i)$
	$P_a \chi_3(k, l)$	ϕ_4	$P_a I_3$	$\frac{q^4 - 3q^2 + 2}{2}$	$c_{7,0}(i, j)$
	$P_a \chi_4(k)$	$\phi_1\phi_2$	$P_a I_4$	$\frac{q^4 - q^2}{2}$	$c_{2,0}(i)$
	$P_a \chi_5(k)$	$\phi_1\phi_2\phi_4$	$P_a I_5$	$q^2 - 1$	$c_{3,0}(i)$
	$P_a \chi_6$	$\phi_1^2\phi_2^2\phi_4$	$P_a I_6$	1	
P_b	$P_b \chi_1(k)$	1	$P_b I_1$	$q^2 - 1$	$c_{2,0}(i)$
	$P_b \chi_5(k, l)$	ϕ_8	$P_b I_5$	$\frac{1}{2}(q^4 - 3q^2 + 2)$	$c_{7,0}(i, j)$
	$P_b \chi_6(k)$	$\phi_1\phi_2\phi_8''$	$P_b I_6$	$\frac{1}{4}(q^4 + \sqrt{2}q^3 - q^2 - \sqrt{2}q)$	$c_{11,0}(i)$
	$P_b \chi_7(k)$	$\phi_1\phi_2\phi_8'$	$P_b I_7$	$\frac{1}{4}(q^4 - \sqrt{2}q^3 - q^2 + \sqrt{2}q)$	$c_{9,0}(i)$
	$P_b \chi_8(k)$	$\phi_1\phi_2\phi_8$	$P_b I_8$	$q^2 - 1$	$c_{3,0}(i)$
	$P_b \chi_{11}$	$\phi_1^2\phi_2^2\phi_8$	$P_b I_{11}$	1	

Table A.16 *Number of fixed points of $H = \langle \alpha^t \rangle$ on parameter sets of the irreducible characters. The unions of parameter sets in this table are disjoint unions.*

Parameter set I	Number of fixed points $ C_I(H) $
GI^{ss}	2^{2t}
$GI_2 \cup GI_3$	2
$GI_4 \cup GI_{30}$	2
$GI_5 \cup GI_6 \cup GI_8 \cup GI_9 \cup GI_{11} \cup GI_{12} \cup GI_{13} \cup GI_{14}$	8
$GI_7 \cup GI_{10}$	2
GI_{17}	1
GI_{18}	1
$GI_{19} \cup GI_{20}$	2
$GI_{23} \cup GI_{43} \cup GI_{47}$	$2^t - 1$
$GI_{24} \cup GI_{44} \cup GI_{48}$	$2^t - 1$
$GI_{25} \cup GI_{45} \cup GI_{49}$	$2^t - 1$
$GI_{27} \cup GI_{33}$	$2^t - 2$
GI_{31}	1
BI_1	$(2^t - 1)^2$
BI_2	$2^t - 1$
BI_3	$2^t - 1$
BI_4	$2^t - 1$
BI_5	$2^t - 1$
$BI_6 \cup BI_7$	2
BI_8	1
BI_9	$2^t - 1$
BI_{10}	1
BI_{11}	$2^t - 1$
BI_{12}	$2^t - 1$
$BI_{13} \cup BI_{14}$	2
$BI_{15} \cup BI_{16} \cup BI_{17} \cup BI_{18} \cup BI_{19} \cup BI_{20} \cup BI_{21} \cup BI_{22}$	8
BI_{23}	$2^t - 1$
BI_{26}	$2^t - 1$
BI_{27}	$2^t - 1$
$BI_{28} \cup BI_{29}$	2
BI_{39}	$2^t - 1$
BI_{42}	$2^t - 1$
BI_{43}	1
BI_{44}	$2^t - 1$
BI_{45}	$2^t - 1$
BI_{51}	$2^t - 1$
BI_{52}	$2^t - 1$
$BI_{53} \cup BI_{54}$	2
$BI_{55} \cup BI_{56} \cup BI_{57} \cup BI_{58}$	4

Table A.16 (cont.) *Number of fixed points of $H = \langle \alpha^t \rangle$ on parameter sets of the irreducible characters.*

Parameter set I	Number of fixed points $ C_I(H) $
$P_a I_1$	$2^t - 1$
$P_a I_2$	$2^t - 1$
$P_a I_3 \cup P_a I_4$	$(2^t - 1)^2$
$P_a I_5$	$2^t - 1$
$P_a I_6$	1
$P_a I_7$	$2^t - 1$
$P_a I_8$	$2^t - 1$
$P_a I_9 \cup P_a I_{10}$	2
$P_a I_{11}$	1
$P_a I_{12}$	$2^t - 1$
$P_a I_{13} \cup P_a I_{25}$	2
$P_a I_{14} \cup P_a I_{15} \cup P_a I_{16} \cup P_a I_{17} \cup P_a I_{21} \cup P_a I_{22} \cup P_a I_{23} \cup P_a I_{24}$	8
$P_a I_{20}$	1
$P_a I_{26}$	$2^t - 1$
$P_a I_{29}$	1
$P_a I_{30}$	1
$P_a I_{31}$	1
$P_a I_{32} \cup P_a I_{33}$	$2^t - 1$
$P_a I_{34}$	$2^t - 1$
$P_a I_{35}$	$2^t - 1$
$P_b I_1$	$2^t - 1$
$P_b I_2$	$2^t - 1$
$P_b I_3$	$2^t - 1$
$P_b I_4$	$2^t - 1$
$P_b I_5 \cup P_b I_6 \cup P_b I_7$	$(2^t - 1)^2$
$P_b I_8$	$2^t - 1$
$P_b I_9 \cup P_b I_{10}$	2
$P_b I_{11}$	1
$P_b I_{12}$	$2^t - 1$
$P_b I_{13}$	1
$P_b I_{14} \cup P_b I_{15}$	2
$P_b I_{16} \cup P_b I_{21} \cup P_b I_{26}$	$2^t - 1$
$P_b I_{17} \cup P_b I_{18} \cup P_b I_{19} \cup P_b I_{20} \cup P_b I_{22} \cup P_b I_{23} \cup P_b I_{24} \cup P_b I_{25}$	8
$P_b I_{27}$	$2^t - 1$
$P_b I_{28}$	$2^t - 1$
$P_b I_{29} \cup P_b I_{30}$	2
$P_b I_{40}$	$2^t - 1$
$P_b I_{43} \cup P_b I_{50}$	2
$P_b I_{44} \cup P_b I_{45} \cup P_b I_{51} \cup P_b I_{52}$	4
$P_b I_{46} \cup P_b I_{53}$	2
$P_b I_{47} \cup P_b I_{48} \cup P_b I_{49}$	$2^t - 1$
$P_b I_{54} \cup P_b I_{55} \cup P_b I_{56}$	$2^t - 1$

APPENDIX B


The irreducible characters of the simple Ree groups ${}^2F_4(q^2)$ were computed by G. Malle, see [25] and the CHEVIE library [14]. However, it seems that 10 families of irreducible characters of ${}^2F_4(q^2)$ have never been published. We give a brief sketch of the construction of these irreducible characters using techniques from Deligne-Lusztig theory. We proceed very much along the same line as F. Lübeck [24]. For many of our calculations we used CHEVIE and GAP [13]. We have implemented the completed character table of ${}^2F_4(q^2)$ as a generic character table in CHEVIE, and whenever we write ${}_G\chi_i(k, \dots)$ we mean the character of ${}^2F_4(q^2)$ which is labeled by $\chi_i(k, \dots)$ in this CHEVIE table. The tables in this Appendix might also be of independent interest; Tables B.9 and B.10 are also used in [15].

Notation and setup. We fix an integer $n > 0$ and set $\theta = 2^n$ and $q := \sqrt{2^{2n+1}} \in \mathbb{R}_{>0}$. Let $\mathbb{F}_{q^2}$ be a finite field with q^2 elements, $\mathbb{F}$ its algebraic closure. To describe our calculations, we fix a simple algebraic group $\mathbf{G}$ of adjoint type F_4 defined over $\mathbb{F}$. Up to isomorphism of algebraic groups, $\mathbf{G}$ is determined by a root datum $(X, \Phi, Y, \Phi^\vee)$, see Section 1.9 in [7]:

Let $X = Y = \mathbb{Z}^4$ be free abelian groups of rank 4 and let $\langle \cdot, \cdot \rangle : X \times Y \rightarrow \mathbb{Z}$, $(x, y) \mapsto xy^{tr}$ be the natural pairing. Let $\{\hat{e}_1, \hat{e}_2, \hat{e}_3, \hat{e}_4\}$ and $\{e_1, e_2, e_3, e_4\}$ be the standard bases of X and Y , respectively. We can describe the root system Φ and the coroot system $\Phi^\vee$ by giving a set $\Delta = \{r_1, r_2, r_3, r_4\} \subseteq X$ of simple roots and a set $\Delta^\vee = \{r_1^\vee, r_2^\vee, r_3^\vee, r_4^\vee\} \subseteq Y$ of simple coroots. We write r_1, r_2, r_3, r_4 as rows of a matrix A and $r_1^\vee, r_2^\vee, r_3^\vee, r_4^\vee$ as rows of a matrix $A^\vee$:

$$A = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \text{and} \quad A^\vee = \begin{pmatrix} 2 & -1 & 0 & 0 \\ -1 & 2 & -1 & 0 \\ 0 & -2 & 2 & -1 \\ 0 & 0 & -1 & 2 \end{pmatrix}.$$

So, the Cartan matrix $\langle r_i, r_j^\vee \rangle$ is equal to $A^\vee$ and we see that the Dynkin diagram of Φ is



To define a Frobenius morphism $F : \mathbf{G} \rightarrow \mathbf{G}$ it is enough to describe the induced action of F on X in the form $q \cdot F_0$ where q is as above and F_0 is the automorphism of the vector space $V = X \otimes_{\mathbb{Z}} \mathbb{R}$ given by $(x_1, x_2, x_3, x_4) \mapsto (\frac{x_4}{\sqrt{2}}, \frac{x_3}{\sqrt{2}}, \frac{2x_2}{\sqrt{2}}, \frac{2x_1}{\sqrt{2}})$. This determines ${}^2F_4(q^2) := \mathbf{G}^F$ up to isomorphism.

A realization of the group $\mathbf{G}$ with Frobenius map F is described in [17] and [28] (and originally in [27]): Let V be a Euclidean vector space with scalar product $(\cdot, \cdot)$, let $\{\varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_4\}$ be an orthonormal basis of V and let Φ be the set consisting of the 48 vectors

$$\varepsilon_i, \varepsilon_i + \varepsilon_j, \frac{1}{2}(\varepsilon_i + \varepsilon_j + \varepsilon_k + \varepsilon_l),$$

where $i, j, k, l \in \{\pm 1, \pm 2, \pm 3, \pm 4\}$, $|i|, |j|, |k|, |l|$ are different and $\varepsilon_{-i} = -\varepsilon_i$ for all i . The set Φ is a root system of type F_4 and the set $\Delta := \{r_1, r_2, r_3, r_4\}$ with the simple roots $r_1 := \varepsilon_2 - \varepsilon_3$, $r_2 := \varepsilon_3 - \varepsilon_4$, $r_3 := \varepsilon_4$, $r_4 := \frac{1}{2}(\varepsilon_1 - \varepsilon_2 - \varepsilon_3 - \varepsilon_4)$ is a basis of Φ . For each $r \in \Phi$, we define the corresponding coroot $r^\vee := \frac{2r}{(r, r)}$. Then, $\Phi^\vee := \{r^\vee \mid r \in \Phi\}$ is a root system of type F_4 with basis $\{r_1^\vee, r_2^\vee, r_3^\vee, r_4^\vee\}$. The

Weyl group $\mathbf{W}$ is the group generated by $w_{r_1}, w_{r_2}, w_{r_3}, w_{r_4}$, the reflections at the hyperplane orthogonal to the simple roots.

Let $\mathbf{G}$ be the simple algebraic group over $\mathbb{F}$ of adjoint type with Frobenius morphism $F : \mathbf{G} \rightarrow \mathbf{G}$, maximal F -stable torus $\mathbf{T}$ and root system Φ which is described in [17] and [28]. In [7, Section 1.9] it is shown how the character group X and the cocharacter group Y of the torus $\mathbf{T}$ are endowed with a natural pairing $\langle \cdot, \cdot \rangle$ and how Φ and $\Phi^\vee$ can be naturally embedded into X and Y , respectively. This gives a root datum $(X, \Phi, Y, \Phi^\vee)$ with the properties described at the beginning of this section.

The elements of the maximal torus $\mathbf{T}$ can be parameterized as follows: There is a natural isomorphism between the abelian groups $\mathbf{T}$ and $\text{Hom}(X, \mathbb{F}^\times)$ (see [7, Section 1.11 and Proposition 3.1.2 (i)]) and we write $h(z_1, z_2, z_3, z_4)$ for the element of $\mathbf{T}$ corresponding to $\chi \in \text{Hom}(X, \mathbb{F}^\times)$ with $\chi(\varepsilon_i) = z_i$ ($i = 1, 2, 3, 4$).

We mention that there is an alternative parameterization of the elements of $\mathbf{T}$ (which in fact we use for our GAP-programs): By [7, Proposition 3.1.2 (ii)], we have $\mathbf{T} \cong Y \otimes \mathbb{F}^\times$ as abelian groups. Every element of $Y \otimes \mathbb{F}^\times$ can be written uniquely as $\sum_{i=1}^4 e_i \otimes \lambda_i$ with $\lambda_i \in \mathbb{F}^\times$ and we write $(\lambda_1, \lambda_2, \lambda_3, \lambda_4)$ for the corresponding element of $\mathbf{T}$. The maps transforming one parameterization into the other are given by:

$$\begin{aligned} (\lambda_1, \lambda_2, \lambda_3, \lambda_4) &\mapsto h(\lambda_1 \lambda_2^2 \lambda_3^3 \lambda_4^2, \lambda_1 \lambda_2 \lambda_3, \lambda_2 \lambda_3, \lambda_3) \quad \text{and} \\ h(z_1, z_2, z_3, z_4) &\mapsto (z_2 z_3^{-1}, z_3 z_4^{-1}, z_4, (z_1 z_2^{-1} z_3^{-1} z_4^{-1})^{1/2}). \end{aligned}$$

The Weyl group $\mathbf{W}$ is isomorphic with $N_{\mathbf{G}}(\mathbf{T})/\mathbf{T}$ and so acts on $\mathbf{T}$ by conjugation which also induces an action of $\mathbf{W}$ on X and on Y , see [24]. The action of $\mathbf{W}$ on X and Y can be determined by the formulas in [7, p. 19]. Once we know the action of $\mathbf{W}$ on X , we can also describe the action on $\mathbf{T}$ via $\mathbf{T} \cong \text{Hom}(X, \mathbb{F}^\times)$. These actions are given in Table B.1.

Finally, we mention that [17, Section 2] provides generators and relations for the group $\mathbf{G}$. Using GAP programs written by Christoph Köhler and the first author in the language of the CHEVIE package, this enables us to carry out explicit computations in the groups $\mathbf{G}$ and $\mathbf{G}^F = {}^2F_4(q^2)$.

Maximal tori. Every semisimple element $s \in \mathbf{G}^F$ is contained in an F -stable maximal torus $\tilde{\mathbf{T}}$ of $\mathbf{G}$, see [12, Corollary 3.16]. Suppose $\tilde{\mathbf{T}}$ is such an F -stable maximal torus of $\mathbf{G}$. We choose some $g \in \mathbf{G}$ such that $\tilde{\mathbf{T}}^g = \mathbf{T}$ and we have $n := g^{-1}F(g) \in N_G(\mathbf{T})$. The action of F on $\tilde{\mathbf{T}}$ corresponds to the action of $(Fn^{-1}) : \mathbf{T} \rightarrow \mathbf{T}, t \mapsto {}^nF(t)$ (in the sense $F(\tilde{t})^g = {}^nF(\tilde{t}^g)$). Since $\mathbf{T}$ is abelian we also write $(Fw^{-1}) : \mathbf{T} \rightarrow \mathbf{T}, t \mapsto {}^wF(t)$ where $w := n\mathbf{T} \in \mathbf{W}$. The map $\tilde{\mathbf{T}} \mapsto w := n\mathbf{T}$ induces a bijection between the $\mathbf{G}^F$ -conjugacy classes of the F -stable maximal tori of $\mathbf{G}$ and the F -conjugacy classes of the Weyl group $\mathbf{W}$, for details see [7, Section 3.3]. A set of representatives w_i for the F -conjugacy classes of $\mathbf{W}$ and the corresponding subgroups $T^{Fw_i^{-1}}$ of fixed points were computed in [28, §3]. A list of these representatives, the order of the corresponding F -centralizers and the maximal tori $T^{(Fw_i^{-1})}$ are given in Table B.2. Parameters for the elements of $T^{(Fw_i^{-1})}$ can be found in Table B.3, where we describe field elements using the notation in [17, Table 5].

Let $(X, \Phi, Y, \Phi^\vee)$ be the root datum of $\mathbf{G}$ described at the beginning of this appendix. So the dual group $\mathbf{G}^*$ is a connected reductive group with root datum

$(Y, \Phi^\vee, X, \Phi)$. The action of F on X and Y gives rise to a Frobenius morphism $F^* : \mathbf{G}^* \rightarrow \mathbf{G}^*$ which determines the finite group $\mathbf{G}^{*F^*}$ up to isomorphism.

A maximal F^* -stable torus of $\mathbf{G}^*$ is isomorphic with $X \otimes_{\mathbb{Z}} \mathbb{F}^\times$ and, via the isomorphism φ_1 from [17, Section 2], also isomorphic with $X \otimes_{\mathbb{Z}} \mathbb{Q}_{p'}/\mathbb{Z}$. Using the $\mathbb{Z}$ -basis $\{\hat{e}_1, \hat{e}_2, \hat{e}_3, \hat{e}_4\}$ of X , every element of $X \otimes_{\mathbb{Z}} \mathbb{Q}_{p'}/\mathbb{Z}$ can be written as

$$s(\mu_1, \mu_2, \mu_3, \mu_4) := \sum_{i=1}^4 \hat{e}_i \otimes \mu_i \text{ with } \mu_i \in \mathbb{Q}_{p'}/\mathbb{Z}.$$

The action of F^* on $\mathbf{T}^*$ is induced by the action of F on X

$$F^*(s(\mu_1, \mu_2, \mu_3, \mu_4)) = s(\theta\mu_4, \theta\mu_3, 2\theta\mu_2, 2\theta\mu_1).$$

Similarly, the action of $\mathbf{W}$ on X induces an action of $\mathbf{W}$ on $\mathbf{T}^*$. This enables us to compute the fixed point groups $\mathbf{T}_i^{*F^*} := (X \otimes_{\mathbb{Z}} \mathbb{Q}_{p'}/\mathbb{Z})^{(w_i F)}$, where w_i runs through the representatives w_i for the F -conjugacy classes of $\mathbf{W}$. In this way, we get the parameterization of the maximal tori of $\mathbf{G}^{*F^*}$ which is described in Table B.4.

For each i , the maximal torus $\mathbf{T}_i^{*F^*}$ of the dual group is isomorphic with the group of linear characters of the torus $\mathbf{T}_i^F$ of $\mathbf{G}^F$ and this isomorphism can be calculated explicitly as in [24, p. 23]. Applying this isomorphism we get the results in Table B.5. These linear characters will be used to parameterize the Lusztig series of irreducible characters of $\mathbf{G}^F$.

Conjugacy classes. We say that two semisimple elements $s_1, s_2 \in \mathbf{G}^F$ are in the same *class type* if their centralizers $C_{\mathbf{G}}(s_1)$ and $C_{\mathbf{G}}(s_2)$ are $\mathbf{G}^F$ -conjugate. A subset $\Psi \subseteq \Phi$ is called a *closed subsystem* if $\mathbb{Z}\Psi \cap \Phi = \Psi$. Let M be the set of all pairs (Π, w) such that Π is a set of simple roots for a closed subsystem of Φ and w is an element of $\mathbf{W}$ such that Π is (Fw^{-1}) -invariant. We call $(\Pi_1, w_1), (\Pi_2, w_2) \in M$ equivalent if and only if there is some $v \in \mathbf{W}$ such that $\Pi_1 v = \Pi_2$ and $v^{-1}w_1 F(v) = w_2$ and let $\bar{M}$ be the set of equivalence classes. The semisimple class types of $\mathbf{G}^F$ can be parameterized by a subset of $\bar{M}$, see [23, Section III.4].

A set of representatives for the semisimple conjugacy classes of $\mathbf{G}^F$ was computed by K. Shinoda, see [28, Table IV], and is given in Table B.7. The semisimple class types of $\mathbf{G}^F$ are listed in Table B.6. The data in these tables can be obtained by the methods described in [23, Section III.4] and [24, Section 4.1]. The notation for the positive roots r_i in Table B.6 is defined in [17, Section 2]. Analogously, we can compute the semisimple class types and representatives for the semisimple conjugacy classes of $\mathbf{G}^{*F^*}$ given in Tables B.9 and B.10. We just have to use the dual torus $\mathbf{T}^*$ instead of $\mathbf{T}$ and the coroots instead of the roots.

Representatives for the unipotent and mixed conjugacy classes were calculated by K. Shinoda, see [28, Tables II and V] and are given in Table B.8. They can be obtained by explicit computations in $\mathbf{G}^F$ using the relations in [17, Tables 1-4].

Lusztig series and Deligne-Lusztig characters. Let $\mathbf{T}_i$ be one of the F -stable maximal tori of $\mathbf{G}$ in Table B.3 and θ a linear character of $\mathbf{T}_i^F$. Associated with the pair $(\mathbf{T}_i, \theta)$ is a generalized character $R_{\mathbf{T}_i}^{\mathbf{G}} \theta$ of $\mathbf{G}^F$, called *Deligne-Lusztig character*. In fact, $R_{\mathbf{T}_i}^{\mathbf{G}} \theta$ only depends on the $\mathbf{G}^F$ -conjugacy class of $(\mathbf{T}_i, \theta)$. Every irreducible character of $\mathbf{G}^F$ occurs as a constituent of some $R_{\mathbf{T}_i}^{\mathbf{G}} \theta$. Every pair $(\mathbf{T}_i, \theta)$ can be identified with a pair $(\mathbf{T}_i^*, s)$ where $\mathbf{T}_i^*$ is one of the F^* -stable maximal tori of the dual group $\mathbf{G}^*$ in Table B.4 and $s \in \mathbf{T}_i^{*F^*}$. Two Deligne-Lusztig characters $R_{\mathbf{T}_i}^{\mathbf{G}} \theta_1$

and $R_{\mathbf{T}_j}^{\mathbf{G}} \theta_2$ can have a common constituent only if the corresponding semisimple elements $s_1, s_2 \in \mathbf{G}^{*F*}$ are $\mathbf{G}^{*F*}$ -conjugate (here we use that the centralizers $C_{\mathbf{G}^*}(s_1), C_{\mathbf{G}^*}(s_2)$ are connected). This induces an equivalence relation on $\text{Irr}(\mathbf{G}^F)$ in a natural way. The equivalence classes $\mathcal{E}(\mathbf{G}^F, (s))$ are parameterized by the semisimple conjugacy classes (s) of $\mathbf{G}^{*F*}$ and are called *Lusztig series*. We say, that a class function is contained in the Lusztig series $\mathcal{E}(\mathbf{G}^F, (s))$ if and only if it is in the $\mathbb{C}$ -span of the irreducible characters in $\mathcal{E}(\mathbf{G}^F, (s))$.

We say that two Lusztig series $\mathcal{E}(\mathbf{G}^F, (s_1)), \mathcal{E}(\mathbf{G}^F, (s_2))$ are of the same type if and only if s_1, s_2 belong to the same semisimple class type of $\mathbf{G}^{*F*}$. So, Tables B.9 and B.10 give an overview over the Lusztig series of $\mathbf{G}^F$. The irreducible characters of $\mathbf{G}^F$ we want to construct are those in the Lusztig series of types $g_8, g_{10}, g_{13}, g_{14}$.

The values of the Deligne-Lusztig characters can be computed using the formula given in [24, Satz 2.1 (b)]: Fix an F -stable maximal torus $\mathbf{T}_i$ of $\mathbf{G}$ and a linear character θ of $\mathbf{T}_i^F$. Let g be an element of $\mathbf{G}^F$ and $g = su = us$ with $s \in \mathbf{G}^F$ semisimple and $u \in \mathbf{G}^F$ unipotent be the Jordan decomposition of g . We write $\mathbf{C} := C_{\mathbf{G}}(s)$ for the centralizer of g in $\mathbf{G}$. Note that in our situation, $\mathbf{C}$ is connected (because $\mathbf{G}$ is simply connected). Let $\mathbf{T}_{i1}, \dots, \mathbf{T}_{ir}$ be a set of representatives for the $\mathbf{C}^F$ -conjugacy classes of F -stable maximal tori in $\mathbf{C}$ which are contained in the $\mathbf{G}^F$ -conjugacy class of $\mathbf{T}_i$ and let $s_{ir1}, \dots, s_{irl_r}$ for $1 \leq r \leq k$ be the different elements of the form s^x with $x \in \mathbf{G}^F$ such that ${}^x\mathbf{T}_i$ is in the $\mathbf{C}^F$ -conjugacy class of $\mathbf{T}_{ir}$. Then

$$(2) \quad (R_{\mathbf{T}_i}^{\mathbf{G}} \theta)(g) = \sum_{r=1}^k \left(\sum_{t=1}^{l_r} \theta(s_{irt}) \right) Q_{\mathbf{T}_{ir}}^{\mathbf{C}}(u)$$

where $Q_{\mathbf{T}_{ir}}^{\mathbf{C}}$ is the Green function of $\mathbf{C}$ with respect to $\mathbf{T}_{ir}$. We will only need $R_{\mathbf{T}_i}^{\mathbf{G}} \theta$ for $i = 3, 4, 5, 6, 7$ and arbitrary θ . The values of $R_{\mathbf{T}_i}^{\mathbf{G}} \theta$ for $i = 3, 4, 5$ can be read off from the character table of ${}^2F_4(q^2)$ in CHEVIE, see the characters $-_G\chi_{35}(k)$, $-_G\chi_{36}(k)$, $_G\chi_{37}(k, l)$ in the CHEVIE table.

So we only have to compute the values of $R_{\mathbf{T}_6}^{\mathbf{G}} \theta$ and $R_{\mathbf{T}_7}^{\mathbf{G}} \theta$. For these Deligne-Lusztig characters, the right hand side of (2) can only be non-zero if s is of class type h_1, h_8, h_{10}, h_{13} or h_{14} . For these classes, $\mathbf{C}$ has Dynkin type ${}^2F_4, {}^2B_2$ or A_0 . So the corresponding Green functions are known, see the CHEVIE library of Green functions.

The elements s_{irt} can be computed from the root datum of $\mathbf{G}$ as it is described in [24, p. 29, (3)]. It turns out that in the computations for $R_{\mathbf{T}_6}^{\mathbf{G}} \theta$ and $R_{\mathbf{T}_7}^{\mathbf{G}} \theta$ we always have $r = k = 1$. This is all information which is needed to compute the values of $R_{\mathbf{T}_6}^{\mathbf{G}} \theta$ and $R_{\mathbf{T}_7}^{\mathbf{G}} \theta$ via formula (2).

The Lusztig series g_{13} and g_{14} correspond to regular semisimple elements of $\mathbf{G}^*$. This implies that $R_{\mathbf{T}_6}^{\mathbf{G}} \theta$ and $R_{\mathbf{T}_7}^{\mathbf{G}} \theta$ are (up to possibly a sign) irreducible characters of $\mathbf{G}^F$. Considering the value on the identity element, we see that in our situation this sign is $+1$. This gives us the irreducible characters $_G\chi_{50}(k, l)$ and $_G\chi_{51}(k, l)$ of $\mathbf{G}^F$ and completes the construction of all irreducible characters in the Lusztig series of types g_{13}, g_{14} . However, since some values of these irreducible characters involve sums of up to 96 roots of unity, we cannot print their values here in this article.

Construction of irreducible characters of type g_8 and g_{10} . Our main tool to construct the irreducible characters in the Lusztig series of types g_8 and g_{10}

is Lusztig's Jordan decomposition of characters: Let $\mathbf{T}_i$ be an F -stable maximal torus of $\mathbf{G}$ and θ a linear character of $\mathbf{T}_i^F$ and let $(\mathbf{T}_i^*, s)$ be a pair corresponding to $(\mathbf{T}_i, \theta)$. Then, there is a bijection

$$\mathcal{L}' : \mathcal{E}(\mathbf{G}^F, (s)) \rightarrow \mathcal{E}(C_{\mathbf{G}^*}(s)^{F^*}, (1)).$$

and for $\chi \in \mathcal{E}(\mathbf{G}^F, (s))$ we have the following identity of scalar products:

$$(3) \quad (\chi, R_{\mathbf{T}_i}^{\mathbf{G}} \theta)_{\mathbf{G}^F} = \varepsilon_{\mathbf{G}} \cdot \varepsilon_{C_{\mathbf{G}^*}(s)} \cdot (\mathcal{L}'(\chi), R_{\mathbf{T}_i^*}^{C_{\mathbf{G}^*}(s)^*} \mathbf{1})_{C_{\mathbf{G}^*}(s)^{F^*}}$$

where $\varepsilon_{\mathbf{G}}, \varepsilon_{C_{\mathbf{G}^*}(s)} \in \{1, -1\}$ and $\mathbf{1}$ is the trivial character of $\mathbf{T}_i^{*F^*}$. The signs $\varepsilon_{\mathbf{G}}, \varepsilon_{C_{\mathbf{G}^*}(s)}$ are defined in [7, Section 6.5] and can be computed directly from their definition. Alternatively, since ${}^2F_4(q^2)$ is isomorphic with its dual group, they can also be read off from the values of the Steinberg character, see [7, Theorem 6.5.9].

We describe the construction of the irreducible characters of $\mathbf{G}^F$ in the Lusztig series of type g_8 . The calculations for the series of type g_{10} are analogous. Let $g_8(k)$ be a representative for the semisimple conjugacy classes of $\mathbf{G}^{F^*}$ as in Table B.10 and let $C^* := C_{\mathbf{G}^*}(g_8(k))$. By Table B.9 we know that C^{*F^*} has Dynkin type 2B_2 . So the Jordan decomposition of characters implies that the Lusztig series $\mathcal{E}(\mathbf{G}^F, (g_8(k)))$ contains exactly 4 irreducible characters, one of them corresponding to the trivial character, two corresponding to the cuspidal characters ${}^2B_2[a], {}^2B_2[b]$ and one corresponding to the Steinberg character for groups of type 2B_2 . We denote these characters, in this order, by ${}_{G\chi_{8,1}}(k), {}_{G\chi_{8,2}}(k), {}_{G\chi_{8,3}}(k), {}_{G\chi_{8,4}}(k)$.

As described in [24, Section 4.1 (2), (3) and Section 6] we can compute representatives for the $\mathbf{G}^{*F^*}$ -conjugacy classes $g_8(k)$ in the various tori $\mathbf{T}_i^*$ and see that the only maximal tori containing such representatives are $\mathbf{T}_3^*, \mathbf{T}_5^*$ and $\mathbf{T}_6^*$. We obtain that, for fixed k , the Deligne-Lusztig characters $-{}_{G\chi_{35}}((2\theta^3 - \theta)k), {}_{G\chi_{37}}(0, k), {}_{G\chi_{50}}(0, k)$ all correspond to the $\mathbf{G}^{*F^*}$ -conjugacy class $g_8(k)$. Using CHEVIE, we can easily compute the norm $(-{}_{G\chi_{35}}((2\theta^3 - \theta)k), -{}_{G\chi_{35}}((2\theta^3 - \theta)k)) = 2$ and, using the scalar product property (3), we can compute the scalar products $(-{}_{G\chi_{35}}((2\theta^3 - \theta)k), {}_{G\chi_{8,1}}(k)) = (-{}_{G\chi_{35}}((2\theta^3 - \theta)k), {}_{G\chi_{8,4}}(k)) = -1$. So that we get

$$-{}_{G\chi_{35}}((2\theta^3 - \theta)k) = -{}_{G\chi_{8,1}}(k) - {}_{G\chi_{8,4}}(k).$$

Analogously, we obtain

$$\begin{aligned} {}_{G\chi_{37}}(0, k) &= -{}_{G\chi_{8,1}}(k) - {}_{G\chi_{8,2}}(k) - {}_{G\chi_{8,3}}(k) + {}_{G\chi_{8,4}}(k), \\ {}_{G\chi_{50}}(0, k) &= -{}_{G\chi_{8,1}}(k) + {}_{G\chi_{8,2}}(k) + {}_{G\chi_{8,3}}(k) + {}_{G\chi_{8,4}}(k). \end{aligned}$$

So, we see that the class function

$$f_8(k) := {}_{G\chi_{8,2}}(k) - {}_{G\chi_{8,3}}(k)$$

is orthogonal to the $\mathbb{C}$ -vector space spanned by the Deligne-Lusztig characters. By inverting the coefficient matrix of the above linear equations, we can now write ${}_{G\chi_{8,1}}(k), \dots, {}_{G\chi_{8,4}}(k)$ as linear combinations of the Deligne-Lusztig characters and $f_8(k)$ as follows:

$$\begin{pmatrix} {}_{G\chi_{8,1}}(k) \\ {}_{G\chi_{8,2}}(k) \\ {}_{G\chi_{8,3}}(k) \\ {}_{G\chi_{8,4}}(k) \end{pmatrix} = \frac{1}{4} \begin{pmatrix} -2 & -1 & -1 & 0 \\ 0 & -1 & 1 & 2 \\ 0 & -1 & 1 & -2 \\ -2 & 1 & 1 & 0 \end{pmatrix} \cdot \begin{pmatrix} -{}_{G\chi_{35}}((2\theta^3 - \theta)k) \\ {}_{G\chi_{37}}(0, k) \\ {}_{G\chi_{50}}(0, k) \\ f_8(k) \end{pmatrix}$$

This already gives us all values of the irreducible characters $_{G\chi_{42}}(k) := _{G\chi_{8,1}}(k)$ and $_{G\chi_{45}}(k) := _{G\chi_{8,4}}(k)$. To determine the values of the non-uniform characters $_{G\chi_{43}}(k)$, $_{G\chi_{44}}(k)$ we have to compute the values of the class functions $f_8(k)$.

The class functions $f_8(k)$. To determine the values of the class function $f_8(k)$ we use arguments similar to those in [24, Section 8]. We proceed in several steps.

Step 1: Class functions constructed from Levi subgroups.

Fix a representative $h \in \mathbf{G}^F$ for the semisimple conjugacy classes of type h_8 as in Table B.7 and let $\mathbf{L}$ be its centralizer in $\mathbf{G}$. Since the order of the representative is not divisible by a bad prime for F_4 , the centralizer $\mathbf{L}$ is an F -stable Levi subgroup of $\mathbf{G}$.

Since $\mathbf{L}^F$ has Dynkin type 2B_2 it has two cuspidal unipotent irreducible characters ${}^2B_2[a]$, ${}^2B_2[b]$ and their difference $f := {}^2B_2[a] - {}^2B_2[b]$ is a unipotent class function of $\mathbf{L}^F$ which is orthogonal to all Deligne-Lusztig characters of $\mathbf{L}^F$. Using the notation from Table B.8, the values of f are:

Class representative	$h_8(i)x_8(1)x_{16}(1)x_{21}(1)$	$h_8(i)x_8(1)x_{16}(1)x_{24}(1)$
Value of f	$\sqrt{2}q\varepsilon_4$	$-\sqrt{2}q\varepsilon_4$

where ε_4 is a complex fourth root of unity as in [17, Table 5]. The values of f on the remaining conjugacy classes of $\mathbf{L}^F$ are zero.

Let $\mathbf{L}^* \subseteq \mathbf{G}^*$ be the dual group of $\mathbf{L}$. Then, $\mathbf{L}^*$ is the centralizer of the semisimple elements of $\mathbf{G}^*$ of type g_8 . Let

$$g_8(k) = s \left(0, \frac{(\theta - 1)k}{q^2 - \sqrt{2}q + 1}, \frac{-k}{q^2 - \sqrt{2}q + 1}, 0 \right)$$

be one of these elements. Since $g_8(k)$ is contained in the center of $\mathbf{L}^{F*}$, we can identify it with a linear character $\lambda(k)$ of $\mathbf{L}^F$ in a natural way, see [24, p. 69]. Using Lusztig induction, we can define the following class functions on $\mathbf{G}^F$:

$$\psi(k) := R_{\mathbf{L}}^{\mathbf{G}}(f \cdot \lambda(k)).$$

Step 2: Connection between $f_8(k)$ and $\psi(k)$.

From [24, Lemma 8.1] and [24, Lemma 8.2] we know that $\psi(k)$ is a generalized character of $\mathbf{G}^F$ in the Lusztig series $g_8(k)$ which is orthogonal to all Deligne-Lusztig characters of $\mathbf{G}^F$. Since the space of class functions in the Lusztig series $g_8(k)$ which are orthogonal to all Deligne-Lusztig characters has dimension one, it follows that $\psi(k)$ is a multiple of $f_8(k)$. Because $\psi(k)$ is a generalized character and f_8 has constituents with multiplicity ± 1 we get that $\psi(k)$ is an integer multiple of $f_8(k)$.

Step 3: Many values of $f_8(k)$, $\psi(k)$ are zero.

The dimension of the space of all class functions which are orthogonal to all Deligne-Lusztig characters of $\mathbf{G}^F$ has dimension $q^2 + 9$ since we get 10 linearly independent such class functions from the unipotent characters and $(q^2 - 2)/2$, $(q^2 - \sqrt{2}q)/4$, $(q^2 + \sqrt{2}q)/4$ such class functions from the Lusztig series of types g_2 , g_8 and g_{10} respectively. The same argument as in [24, p. 71] shows that $\psi(k)$ vanishes on all conjugacy classes of $\mathbf{G}^F$ except for possibly the following tuples of classes: $(c_{1,3}, c_{1,4})$, $(c_{1,7}, c_{1,8}, c_{1,9})$, $(c_{1,10}, c_{1,11}, c_{1,12})$, $(c_{1,13}, c_{1,14})$, $(c_{1,15}, c_{1,16}, c_{1,17}, c_{1,18})$, $(c_{2,2}(i), c_{2,3}(i))$, $(c_{5,2}, c_{5,3}, c_{5,4})$, $(c_{8,2}(i), c_{8,3}(i))$, $(c_{10,2}(i), c_{10,3}(i))$. Additionally,

the argument shows that the values of $\psi(k)$ satisfy the following relations:

$$\begin{aligned}
\psi(k)(c_{1,4}) &= -\psi(k)(c_{1,3}) \\
\psi(k)(c_{1,9}) &= -\psi(k)(c_{1,8}) + \psi(k)(c_{1,7}) \\
\psi(k)(c_{1,12}) &= -2\psi(k)(c_{1,10}) - \psi(k)(c_{1,11}) \\
\psi(k)(c_{1,14}) &= -\psi(k)(c_{1,13}) \\
\psi(k)(c_{1,18}) &= -\psi(k)(c_{1,15}) - \psi(k)(c_{1,16}) - \psi(k)(c_{1,17}) \\
\psi(k)(c_{8,3}(i)) &= -\psi(k)(c_{8,2}(i))
\end{aligned}$$

Step 4: Values of the remaining irreducible characters.

We use the formula [12, Proposition 12.2] for Lusztig induction to derive some information on the non-zero values of $\psi(k)$. (Unfortunately, the formula does not give us the values of $\psi(k)$ explicitly, since we do not know the two-parameter Green functions occurring in the right hand side of the formula.)

Firstly, we can conclude that $\psi(k)$ vanishes on the classes $c_{2,2}(i)$, $c_{2,3}(i)$, $c_{5,2}$, $c_{5,3}$, $c_{5,4}$, $c_{10,2}(i)$, $c_{10,3}(i)$ since the semisimple part of the representatives for these classes cannot be conjugate to an element of the center $Z(\mathbf{L}^F)$. Note that by [28, p.13], we have $\mathbf{L}^F = Z(\mathbf{L}^F) \times (\mathbf{L}^F)'$. Secondly, we see that

$$\begin{aligned}
\psi(k)(c_{8,2}(i)) &= x_{8,2}\varepsilon_4(\varphi_8''^{qik/\sqrt{2}} + \varphi_8''^{-qik/\sqrt{2}} + \varphi_8''^{(q/\sqrt{2}-1)ik} + \varphi_8''^{-(q/\sqrt{2}-1)ik}), \\
\psi(k)(c_{8,3}(i)) &= x_{8,3}\varepsilon_4(\varphi_8''^{qik/\sqrt{2}} + \varphi_8''^{-qik/\sqrt{2}} + \varphi_8''^{(q/\sqrt{2}-1)ik} + \varphi_8''^{-(q/\sqrt{2}-1)ik})
\end{aligned}$$

for some rational numbers $x_{8,2}$, $x_{8,3}$ not depending on i, k . Thirdly, the formula implies that the values of $\psi(k)$ on the unipotent conjugacy classes are of the form $\psi(k)(c_{1,r}) = \varepsilon_4 x_{1,r}$ where $x_{1,r}$ are rational integers. Considering the $x_{1,r}$, $x_{8,r}$ as indeterminates and using CHEVIE, we can compute the scalar products of $\psi(k)$ with the unipotent irreducible characters of $\mathbf{G}^F$. This gives us $\psi(k)$ and hence $f_8(k)$ up to a rational multiple. From the condition $(f_8(k), f_8(k)) = 2$ we get $f_8(k)$ up to a sign, see Table B.11. Note, that the sign of $f_8(k)$ only flips the complex-conjugate irreducible characters $\chi_{8,2}(k)$ and $\chi_{8,3}(k)$. This gives us the irreducible characters $\chi_{43}(k)$ and $\chi_{44}(k)$ and thereby completes the determination of the irreducible characters of $\mathbf{G}^F$ in the Lusztig series of type g_8 . The values of the irreducible characters $\chi_{46}(k), \dots, \chi_{49}(k)$ of $\mathbf{G}^F$ in the Lusztig series of type g_{10} can be computed analogously.

Again, for space reasons, we cannot print all values of the irreducible characters $\chi_{42}(k), \dots, \chi_{49}(k)$. Therefore, Table B.12 presents only the degrees and those character values which are used in the verification of Dade's conjecture.

Remark: We have implemented the completed character table of $\mathbf{G}^F = {}^2F_4(q^2)$ as a generic character table in CHEVIE. To test the correctness of the table, we have verified the row and column orthogonality relations. Furthermore, using CHEVIE, we computed the tensor products of all irreducible characters with the unipotent characters and computed the scalar products (which have to be nonnegative integers) of these tensor products with the unipotent characters.

Table B.1 *Description of the generators $w_{r_1}, w_{r_2}, w_{r_3}, w_{r_4}$ of the Weyl group $\mathbf{W}$ of type F_4 . As automorphisms of the character group X and the cocharacter group Y , we describe the generators by their images of the chosen $\mathbb{Z}$ -bases of X and Y , respectively. As elements of $N_{\mathbf{G}}(\mathbf{T})/\mathbf{T}$, we describe them by giving the torus elements $w_{r_i} h(z_1, z_2, z_3, z_4)$.*

$\mathbf{W}$	w_{r_i}	Action of w_{r_i}	Matrix (Row convention)
in $\text{Aut}(X)$	w_{r_1}	$\hat{e}_1 \mapsto -\hat{e}_1,$ $\hat{e}_2 \mapsto \hat{e}_1 + \hat{e}_2,$ $\hat{e}_3 \mapsto \hat{e}_3,$ $\hat{e}_4 \mapsto \hat{e}_4$	$\begin{pmatrix} -1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$
	w_{r_2}	$\hat{e}_1 \mapsto \hat{e}_1 + \hat{e}_2,$ $\hat{e}_2 \mapsto -\hat{e}_2,$ $\hat{e}_3 \mapsto \hat{e}_2 + \hat{e}_3,$ $\hat{e}_4 \mapsto \hat{e}_4$	$\begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$
	w_{r_3}	$\hat{e}_1 \mapsto \hat{e}_1,$ $\hat{e}_2 \mapsto \hat{e}_2 + 2\hat{e}_3,$ $\hat{e}_3 \mapsto -\hat{e}_3,$ $\hat{e}_4 \mapsto \hat{e}_3 + \hat{e}_4$	$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix}$
	w_{r_4}	$\hat{e}_1 \mapsto \hat{e}_1,$ $\hat{e}_2 \mapsto \hat{e}_2,$ $\hat{e}_3 \mapsto \hat{e}_3 + \hat{e}_4,$ $\hat{e}_4 \mapsto -\hat{e}_4$	$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & -1 \end{pmatrix}$
in $\text{Aut}(Y)$	w_{r_1}	$e_1 \mapsto -e_1 + e_2,$ $e_2 \mapsto e_2,$ $e_3 \mapsto e_3,$ $e_4 \mapsto e_4$	$\begin{pmatrix} -1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$
	w_{r_2}	$e_1 \mapsto e_1,$ $e_2 \mapsto e_1 - e_2 + e_3,$ $e_3 \mapsto e_3,$ $e_4 \mapsto e_4$	$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & -1 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$
	w_{r_3}	$e_1 \mapsto e_1,$ $e_2 \mapsto e_2,$ $e_3 \mapsto 2e_2 - e_3 + e_4,$ $e_4 \mapsto e_4$	$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 2 & -1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}$
	w_{r_4}	$e_1 \mapsto e_1,$ $e_2 \mapsto e_2,$ $e_3 \mapsto e_3,$ $e_4 \mapsto e_3 - e_4$	$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & -1 \end{pmatrix}$
in $N_{\mathbf{G}}(\mathbf{T})/\mathbf{T}$	w_{r_1}	$h(z_1, z_3, z_2, z_4)$	
	w_{r_2}	$h(z_1, z_2, z_4, z_3)$	
	w_{r_3}	$h(z_1, z_2, z_3, z_4^{-1})$	
	w_{r_4}	$h(z_1^{\frac{1}{2}} z_2^{\frac{1}{2}} z_3^{\frac{1}{2}} z_4^{\frac{1}{2}}, z_1^{\frac{1}{2}} z_2^{\frac{1}{2}} z_3^{-\frac{1}{2}} z_4^{-\frac{1}{2}}, z_1^{\frac{1}{2}} z_2^{-\frac{1}{2}} z_3^{\frac{1}{2}} z_4^{-\frac{1}{2}}, z_1^{\frac{1}{2}} z_2^{-\frac{1}{2}} z_3^{-\frac{1}{2}} z_4^{\frac{1}{2}})$	

Table B.2 (cf. [28, Table III]) *The F -conjugacy classes of the Weyl group $\mathbf{W}$ of type F_4 . For each class, we give a representative w_i , the order of the F -centralizer $C_{\mathbf{W},F}(w_i)$ and the order of $\mathbf{T}^{Fw_i^{-1}}$.*

$\mathbf{T}_i$	w_i	$ C_{\mathbf{W},F}(w_i) $	$ \mathbf{T}_i^F $
T_1	1	16	$(q^2 - 1)^2$
T_2	w_{r_1}	4	$q^4 - 1$
T_3	w_{r_3}	8	$(q^2 - 1)(q^2 - \sqrt{2}q + 1)$
T_4	$w_{r_2} w_{r_3} w_{r_2}$	8	$(q^2 - 1)(q^2 + \sqrt{2}q + 1)$
T_5	$w_{r_1} w_{r_2} w_{r_1} w_{r_3} w_{r_2} w_{r_1} w_{r_3} w_{r_2}$	16	$q^4 + 1$
T_6	$w_{r_1} w_{r_2} w_{r_3} w_{r_2} w_{r_1} w_{r_3}$	96	$(q^2 - \sqrt{2}q + 1)^2$
T_7	$w_{r_2} w_{r_3} w_{r_2} w_{r_4} w_{r_3} w_{r_2} w_{r_1} w_{r_3} w_{r_2} \cdot$ $w_{r_3} w_{r_4} w_{r_3} w_{r_2} w_{r_1} w_{r_3} w_{r_2} w_{r_3} w_{r_4}$	96	$(q^2 + \sqrt{2}q + 1)^2$
T_8	$w_{r_1} w_{r_2} w_{r_3} w_{r_4} w_{r_3} w_{r_2} w_{r_1} w_{r_3} w_{r_2} \cdot$ $w_{r_4} w_{r_3} w_{r_2}$	48	$(q^2 + 1)^2$
T_9	$w_{r_2} w_{r_3} w_{r_2} w_{r_1}$	6	$q^4 - q^2 + 1$
T_{10}	$w_{r_1} w_{r_2}$	12	$q^4 - \sqrt{2}q^3 + q^2 - \sqrt{2}q + 1$
T_{11}	$w_{r_1} w_{r_4} w_{r_3} w_{r_2} w_{r_1} w_{r_3} w_{r_2} w_{r_4} w_{r_3} w_{r_2}$	12	$q^4 + \sqrt{2}q^3 + q^2 + \sqrt{2}q + 1$

Table B.3 (cf. Shinoda [28, (3.1)]) *Parameters for the elements of the maximal tori*
 $\mathbf{T}_i^F = \mathbf{T}^{(Fw_i^{-1})}$.

Torus	$\mathbf{T}^{(Fw_i^{-1})}$
$\mathbf{T}_1^F$	$\{t_1(i, j) := h(\tilde{\zeta}^i, \tilde{\zeta}^{(2\theta-1)i}, \tilde{\zeta}^j, \tilde{\zeta}^{(2\theta-1)j}) \mid 0 \leq i, j \leq q^2 - 2\}$
$\mathbf{T}_2^F$	$\{t_2(i) := h(\tilde{\zeta}_4^{(4\theta^3+2\theta^2+1)i}, \tilde{\zeta}_4^{(2\theta+2\theta^2-1)i}, \tilde{\zeta}_4^{(2\theta-2\theta^2+1)i}, \tilde{\zeta}_4^{(2\theta^2-4\theta^3+1)i}) \mid$ $0 \leq i \leq q^4 - 2\}$
$\mathbf{T}_3^F$	$\{t_3(i) := h(\tilde{\psi}_8^{(2\theta^2-2\theta+1)i}, \tilde{\psi}_8^{(4\theta^3-6\theta^2+4\theta-1)i}, \tilde{\psi}_8^{(2\theta^2-1)i}, \tilde{\psi}_8^{(-4\theta^4+2\theta^2)i}) \mid$ $0 \leq i \leq q^4 - \sqrt{2}q^3 + \sqrt{2}q - 2\}$
$\mathbf{T}_4^F$	$\{t_4(i) := h(\tilde{\psi}_8^{(2\theta^2+2\theta+1)i}, \tilde{\psi}_8^{(4\theta^3+2\theta^2-1)i}, \tilde{\psi}_8^{(2\theta^2-1)i}, \tilde{\psi}_8^{(-4\theta^4+2\theta^2)i}) \mid$ $0 \leq i \leq q^4 + \sqrt{2}q^3 - \sqrt{2}q - 2\}$
$\mathbf{T}_5^F$	$\{t_5(i, j) := h(\tilde{\varphi}_8^{''i}, \tilde{\varphi}_8^{''-q^2i}, \tilde{\varphi}_8^{'j}, \tilde{\varphi}_8^{'-q^2j}) \mid 0 \leq i \leq q^2 - \sqrt{2}q, 0 \leq j \leq q^2 + \sqrt{2}q\}$
$\mathbf{T}_6^F$	$\{t_6(i, j) := h(\tilde{\varphi}_8^{''i}, \tilde{\varphi}_8^{''-q^2i}, \tilde{\varphi}_8^{''j}, \tilde{\varphi}_8^{''-q^2j}) \mid 0 \leq i, j \leq q^2 - \sqrt{2}q\}$
$\mathbf{T}_7^F$	$\{t_7(i, j) := h(\tilde{\varphi}_8^{'i}, \tilde{\varphi}_8^{'-q^2i}, \tilde{\varphi}_8^{'j}, \tilde{\varphi}_8^{'-q^2j}) \mid 0 \leq i, j \leq q^2 + \sqrt{2}q\}$
$\mathbf{T}_8^F$	$\{t_8(i, j) := h(\tilde{\xi}_2^i, \tilde{\xi}_2^j, \tilde{\xi}_2^{\theta(i+j)}, \tilde{\xi}_2^{\theta(i-j)}) \mid 0 \leq i, j \leq q^2\}$
$\mathbf{T}_9^F$	$\{t_9(i) := h(\tilde{\varphi}_{12}^i, \tilde{\varphi}_{12}^{(-4\theta^3+2\theta-1)i}, \tilde{\varphi}_{12}^{(1-2\theta)i}, \tilde{\varphi}_{12}^{(4\theta^3-4\theta^2+1)i}) \mid 0 \leq i \leq q^4 - q^2\}$
$\mathbf{T}_{10}^F$	$\{t_{10}(i) := h(\tilde{\varphi}_{24}^{''i}, \tilde{\varphi}_{24}^{''(q^2-q^4)i}, \tilde{\varphi}_{24}^{''(2\theta-1)i}, \tilde{\varphi}_{24}^{''(4\theta^3-1)i}) \mid$ $0 \leq i \leq q^4 - \sqrt{2}q^3 + q^2 - \sqrt{2}q\}$
$\mathbf{T}_{11}^F$	$\{t_{11}(i) := h(\tilde{\varphi}_{24}^{'(q^4-q^2)i}, \tilde{\varphi}_{24}^{'i}, \tilde{\varphi}_{24}^{'(2\theta+1)i}, \tilde{\varphi}_{24}^{'(-4\theta^3-1)i}) \mid$ $0 \leq i \leq q^4 + \sqrt{2}q^3 + q^2 + \sqrt{2}q\}$

Table B.4 *Parameters for the elements of the maximal tori $\mathbf{T}_i^{*F*} = (X \otimes_{\mathbb{Z}} \mathbb{Q}_{p'}/\mathbb{Z})^{w_i F}$ of the dual group $\mathbf{G}^{*F*}$. We use the abbreviations $\phi'_{24} := q^4 + \sqrt{2}q^3 + q^2 + \sqrt{2}q + 1$ and $\phi''_{24} := q^4 - \sqrt{2}q^3 + q^2 - \sqrt{2}q + 1$.*

Torus	$(X \otimes_{\mathbb{Z}} \mathbb{Q}_{p'}/\mathbb{Z})^{w_i F}$
$\mathbf{T}_1^{*F*}$	$\{s_1(k, l) := s\left(\frac{(2\theta+1)k+(2\theta+1)l}{q^2-1}, \frac{(4\theta+3)k+(2\theta+1)l}{q^2-1}, \frac{(6\theta+4)k+(2\theta+2)l}{q^2-1}, \frac{(2\theta+2)k+(2\theta+2)l}{q^2-1}\right) \mid$ $0 \leq k, l \leq q^2 - 2\}$
$\mathbf{T}_2^{*F*}$	$\{s_2(k) := s\left(\frac{2\theta^4(2\theta^3+2\theta^2+\theta)k}{q^4-1}, \frac{2\theta^4(4\theta^3+2\theta^2+2\theta+1)k}{q^4-1}, \frac{2\theta^4(4\theta^2+2\theta+4\theta^3+2)k}{q^4-1}, \frac{2\theta^4(2\theta^2+2\theta+1)k}{q^4-1}\right) \mid$ $0 \leq k \leq q^4 - 2\}$
$\mathbf{T}_3^{*F*}$	$\{s_3(k) := s\left(\frac{(6\theta^3-2\theta^2-\theta+2)k}{(q^2-1)(q^2-\sqrt{2}q+1)}, \frac{(10\theta^3-2\theta^2-2\theta+3)k}{(q^2-1)(q^2-\sqrt{2}q+1)}, \frac{(16\theta^3-4\theta^2-4\theta+5)k}{(q^2-1)(q^2-\sqrt{2}q+1)}, \frac{(8\theta^3-2\theta^2-2\theta+3)k}{(q^2-1)(q^2-\sqrt{2}q+1)}\right) \mid$ $0 \leq k \leq q^4 - \sqrt{2}q^3 + \sqrt{2}q - 2\}$
$\mathbf{T}_4^{*F*}$	$\{s_4(k) := s\left(\frac{(2\theta^3+2\theta^2+\theta)k}{(q^2-1)(q^2+\sqrt{2}q+1)}, \frac{(2\theta^3+4\theta^2+2\theta)k}{(q^2-1)(q^2+\sqrt{2}q+1)}, \frac{(4\theta^2+4\theta+1)k}{(q^2-1)(q^2+\sqrt{2}q+1)}, \frac{(2\theta^2+2\theta+1)k}{(q^2-1)(q^2+\sqrt{2}q+1)}\right) \mid$ $0 \leq k \leq q^4 + \sqrt{2}q^3 - \sqrt{2}q - 2\}$
$\mathbf{T}_5^{*F*}$	$\{s_5(k, l) := s\left(\frac{-k}{q^2-\sqrt{2}q+1}, \frac{(\theta-2)k}{q^2-\sqrt{2}q+1} + \frac{(-\theta-1)l}{q^2+\sqrt{2}q+1}, \frac{(2\theta-3)k}{q^2-\sqrt{2}q+1} + \frac{-l}{q^2+\sqrt{2}q+1}, \frac{(2\theta-2)k}{q^2-\sqrt{2}q+1}\right) \mid$ $0 \leq k \leq q^2 - \sqrt{2}q, 0 \leq l \leq q^2 + \sqrt{2}q\}$
$\mathbf{T}_6^{*F*}$	$\{s_6(k, l) := s\left(\frac{-k}{q^2-\sqrt{2}q+1}, \frac{(\theta-2)k+(\theta-1)l}{q^2-\sqrt{2}q+1}, \frac{(2\theta-3)k-l}{q^2-\sqrt{2}q+1}, \frac{(2\theta-2)k}{q^2-\sqrt{2}q+1}\right) \mid 0 \leq k, l \leq q^2 - \sqrt{2}q\}$
$\mathbf{T}_7^{*F*}$	$\{s_7(k, l) := s\left(\frac{-k}{q^2+\sqrt{2}q+1}, \frac{(-\theta-2)k+(-\theta-1)l}{q^2+\sqrt{2}q+1}, \frac{(-2\theta-3)k-l}{q^2+\sqrt{2}q+1}, \frac{(-2\theta-2)k}{q^2+\sqrt{2}q+1}\right) \mid 0 \leq k, l \leq q^2 + \sqrt{2}q\}$
$\mathbf{T}_8^{*F*}$	$\{s_8(k, l) := s\left(\frac{-2k}{q^2+1}, \frac{(2\theta-3)k-l}{q^2+1}, \frac{(2\theta-4)k+(2\theta-2)l}{q^2+1}, \frac{-2k-2l}{q^2+1}\right) \mid 0 \leq k, l \leq q^2\}$
$\mathbf{T}_9^{*F*}$	$\{s_9(k) := s\left(\frac{(2\theta^3-2\theta-1)k}{q^4-q^2+1}, \frac{(-3\theta-2+2\theta^3)k}{q^4-q^2+1}, \frac{(-4\theta-3+4\theta^3)k}{q^4-q^2+1}, \frac{(4\theta^3+2\theta^2-2\theta-2)k}{q^4-q^2+1}\right) \mid$ $0 \leq k \leq q^4 - q^2\}$
$\mathbf{T}_{10}^{*F*}$	$\{s_{10}(k) := s\left(\frac{(2\theta^3-1)k}{\phi''_{24}}, \frac{(6\theta^3-2\theta^2+\theta-2)k}{\phi''_{24}}, \frac{(8\theta^3-2\theta^2+2\theta-3)k}{\phi''_{24}}, \frac{(4\theta^3-2\theta^2+2\theta-2)k}{\phi''_{24}}\right) \mid$ $0 \leq k \leq q^4 - \sqrt{2}q^3 + q^2 - \sqrt{2}q\}$
$\mathbf{T}_{11}^{*F*}$	$\{s_{11}(k) := s\left(\frac{(-2\theta^3-2\theta^2-2\theta-1)k}{\phi_{24}}, \frac{(-3\theta-2\theta^2-1)k}{\phi_{24}}, \frac{(-4\theta-2\theta^2-1)k}{\phi_{24}}, \frac{(-2\theta^2-2\theta)k}{\phi_{24}}\right) \mid$ $0 \leq k \leq q^4 + \sqrt{2}q^3 + q^2 + \sqrt{2}q\}$

Table B.5 *Interpretation of the elements $s_i(k, \dots)$ of the tori of the dual group as linear characters $\theta_i(k, \dots) : \mathbf{T}_i^F \rightarrow \mathbb{Q}_{p'}/\mathbb{Z}$. From those we get complex-valued linear characters of $\mathbf{T}_i^F$ via the embedding $\varphi_2 : \mathbb{Q}_{p'} \hookrightarrow \mathbb{C}^\times$, $\frac{r}{s} \mapsto e^{2\pi i r/s}$.*

Torus	$\theta_i(k, \dots) : \mathbf{T}_i^F \rightarrow \mathbb{Q}_{p'}/\mathbb{Z}$
$\hat{\mathbf{T}}_1^F$	$\theta_1(k, l) : t_1(a, b) \mapsto \frac{(k+l)a + (k-l)b}{q^2 - 1}$
$\hat{\mathbf{T}}_2^F$	$\theta_2(k) : t_2(a) \mapsto \frac{ak}{q^4 - 1}$
$\hat{\mathbf{T}}_3^F$	$\theta_3(k) : t_3(a) \mapsto \frac{ak}{(q^2 - 1)(q^2 - \sqrt{2}q + 1)}$
$\hat{\mathbf{T}}_4^F$	$\theta_4(k) : t_4(a) \mapsto \frac{ak}{(q^2 - 1)(q^2 + \sqrt{2}q + 1)}$
$\hat{\mathbf{T}}_5^F$	$\theta_5(k, l) : t_5(a, b) \mapsto \frac{ak}{q^2 - \sqrt{2}q + 1} + \frac{bl}{q^2 + \sqrt{2}q + 1}$
$\hat{\mathbf{T}}_6^F$	$\theta_6(k, l) : t_6(a, b) \mapsto \frac{ak + bl}{q^2 - \sqrt{2}q + 1}$
$\hat{\mathbf{T}}_7^F$	$\theta_7(k, l) : t_7(a, b) \mapsto \frac{ak + bl}{q^2 + \sqrt{2}q + 1}$
$\hat{\mathbf{T}}_8^F$	$\theta_8(k, l) : t_8(a, b) \mapsto \frac{(k+l)a + (k-l)b}{q^2 + 1}$
$\hat{\mathbf{T}}_9^F$	$\theta_9(k) : t_9(a) \mapsto \frac{ak}{q^4 - q^2 + 1}$
$\hat{\mathbf{T}}_{10}^F$	$\theta_{10}(k) : t_{10}(a) \mapsto \frac{ak}{q^4 - \sqrt{2}q^3 + q^2 - \sqrt{2}q + 1}$
$\hat{\mathbf{T}}_{11}^F$	$\theta_{11}(k) : t_{11}(a) \mapsto \frac{ak}{q^4 + \sqrt{2}q^3 + q^2 + \sqrt{2}q + 1}$

Table B.6 *Parameterization of the semisimple class types of ${}^2F_4(q^2)$.*

No.	Dynkin type	Π	w
h_1	2F_4	$[r_1, r_2, r_3, r_4]$	1
h_2	2B_2	$[r_8, r_{16}]$	1
h_3	A_1	$[r_{15}, r_{20}]$	1
h_4	A_0	$[\]$	1
h_5	2A_2	if $\theta \equiv 1 \pmod 3$: $[r_8, r_9, r_{10}, r_{18}]$ if $\theta \equiv -1 \pmod 3$: $[r_6, r_{11}, r_{12}, r_{16}]$	w_{r_1}
h_6	A_1	$[r_{17}, r_{22}]$	w_{r_1}
h_7	A_0	$[\]$	w_{r_1}
h_8	2B_2	$[r_8, r_{16}]$	w_{r_3}
h_9	A_0	$[\]$	w_{r_3}
h_{10}	2B_2	$[r_8, r_{16}]$	$w_{r_2} w_{r_3} w_{r_2}$
h_{11}	A_0	$[\]$	$w_{r_2} w_{r_3} w_{r_2}$
h_{12}	A_0	$[\]$	$w_{r_1} w_{r_2} w_{r_1} w_{r_3} w_{r_2} w_{r_1} w_{r_3} w_{r_2}$
h_{13}	A_0	$[\]$	$w_{r_1} w_{r_2} w_{r_3} w_{r_2} w_{r_1} w_{r_3}$
h_{14}	A_0	$[\]$	$w_{r_2} w_{r_3} w_{r_2} w_{r_4} (w_{r_3} w_{r_2} w_{r_1} w_{r_3} w_{r_2} w_{r_3} w_{r_4})^2$
h_{15}	A_0	$[\]$	$w_{r_1} w_{r_2} w_{r_3} w_{r_4} w_{r_3} w_{r_2} w_{r_1} w_{r_3} w_{r_2} w_{r_4} w_{r_3} w_{r_2}$
h_{16}	A_0	$[\]$	$w_{r_2} w_{r_3} w_{r_2} w_{r_1}$
h_{17}	A_0	$[\]$	$w_{r_1} w_{r_2}$
h_{18}	A_0	$[\]$	$w_{r_1} w_{r_4} w_{r_3} w_{r_2} w_{r_1} w_{r_3} w_{r_2} w_{r_4} w_{r_3} w_{r_2}$

Table B.7 (cf. Shinoda [28, Table IV] and [17, Table A.1]) *Parameterization of the semisimple conjugacy classes of ${}^2F_4(q^2)$.*

Representative	Parameters	Number of Classes
$h_1 := h(1, 1, 1, 1)$		1
$h_2(i) := h(1, 1, \tilde{\zeta}_2^i, \tilde{\zeta}_2^{(2\theta-1)i})$	$i = 0, \dots, q^2 - 2$ $i \neq 0$	$\frac{q^2-2}{2}$
$h_3(i) := h(\tilde{\zeta}_2^i, \tilde{\zeta}_2^{(2\theta-1)i}, \tilde{\zeta}_2^{(2\theta+1)i}, \tilde{\zeta}_2^i)$	$i = 0, \dots, q^2 - 2$ $i \neq 0$	$\frac{q^2-2}{2}$
$h_4(i, j) := h(\tilde{\zeta}_2^i, \tilde{\zeta}_2^{(2\theta-1)i}, \tilde{\zeta}_2^j, \tilde{\zeta}_2^{(2\theta-1)j})$	$i = 0, \dots, q^2 - 2; j = 0, \dots, q^2 - 2$ $i \neq 0, \pm j, \pm(2\theta-1)j$ $j \neq 0, \pm i, \pm(2\theta-1)i$	$\frac{q^4-10q^2+16}{16}$
$h_5 := h(\tilde{\varepsilon}_3, \tilde{\varepsilon}_3^{1-\theta}, \tilde{\varepsilon}_3^{-1-\theta}, \tilde{\varepsilon}_3^{-\theta})$		1
$h_6(i) := h(\tilde{\xi}_2^i, \tilde{\xi}_2^{(1-\theta)i}, \tilde{\xi}_2^{(-1-\theta)i}, \tilde{\xi}_2^{-\theta i})$	$i = 0, \dots, q^2$ $i \neq 0, \frac{q^2+1}{3}, \frac{2(q^2+1)}{3}$	$\frac{q^2-2}{2}$
$h_7(i) := h(\tilde{\zeta}_4^{(4\theta^3+2\theta^2+1)i}, \tilde{\zeta}_4^{(2\theta^2+2\theta-1)i}, \tilde{\zeta}_4^{(-2\theta^2+2\theta+1)i}, \tilde{\zeta}_4^{-4\theta^3+2\theta^2+1)i})$	$i = 0, \dots, q^4 - 2$ $i \neq (q^2-1)m, m = 0, \dots, q^2$ $i \neq (q^2+1)m, m = 0, \dots, q^2 - 2$	$\frac{q^4-2q^2}{4}$
$h_8(i) := h(1, 1, \tilde{\varphi}_8^{i'}, \tilde{\varphi}_8^{i'-q^2 i})$	$i = 0, \dots, q^2 - \sqrt{2}q$ $i \neq 0$	$\frac{q^2-\sqrt{2}q}{4}$
$h_9(i) := h(\tilde{\psi}_8^{i'(2\theta^2-2\theta+1)i}, \tilde{\psi}_8^{i'(4\theta^3-6\theta^2+4\theta-1)i}, \tilde{\psi}_8^{i'(2\theta^2-1)i}, \tilde{\psi}_8^{i'(-4\theta^4+2\theta^2)i})$	$i = 0, \dots, q^4 - \sqrt{2}q^3 + \sqrt{2}q - 2$ $i \neq (q^2-1)m, m = 0, \dots, q^2 - \sqrt{2}q$ $i \neq (q^2 - \sqrt{2}q + 1)m, m = 0, \dots, q^2 - 2$	$\frac{1}{8}(q^4 - \sqrt{2}q^3 - 2q^2 + 2\sqrt{2}q)$
$h_{10}(i) := h(1, 1, \tilde{\varphi}_8^{i'}, \tilde{\varphi}_8^{i'-q^2 i})$	$i = 0, \dots, q^2 + \sqrt{2}q$ $i \neq 0$	$\frac{q^2+\sqrt{2}q}{4}$
$h_{11}(i) := h(\tilde{\psi}_8^{i'(2\theta^2+2\theta+1)i}, \tilde{\psi}_8^{i'(4\theta^3+2\theta^2-1)i}, \tilde{\psi}_8^{i'(2\theta^2-1)i}, \tilde{\psi}_8^{i'(-4\theta^4+2\theta^2)i})$	$i = 0, \dots, q^4 + \sqrt{2}q^3 - \sqrt{2}q - 2$ $i \neq (q^2-1)m, m = 0, \dots, q^2 + \sqrt{2}q$ $i \neq (q^2 + \sqrt{2}q + 1)m, m = 0, \dots, q^2 - 2$	$\frac{1}{8}(q^4 + \sqrt{2}q^3 - 2q^2 - 2\sqrt{2}q)$
$h_{12}(i, j) := h(\tilde{\varphi}_8^{i'}, \tilde{\varphi}_8^{i'-q^2 i}, \tilde{\varphi}_8^{j'}, \tilde{\varphi}_8^{j'-q^2 j})$	$i = 0, \dots, q^2 - \sqrt{2}q; j = 0, \dots, q^2 + \sqrt{2}q$ $i, j \neq 0$	$\frac{q^2(q^2-2)}{16}$
$h_{13}(i, j) := h(\tilde{\varphi}_8^{i'}, \tilde{\varphi}_8^{i'-q^2 i}, \tilde{\varphi}_8^{j'}, \tilde{\varphi}_8^{j'-q^2 j})$	$i = 0, \dots, q^2 - \sqrt{2}q; j = 0, \dots, q^2 - \sqrt{2}q$ $i \neq 0, \pm j, \pm q^2 j$ $j \neq 0, \pm i, \pm q^2 i$	$\frac{1}{96}(q^4 - 2\sqrt{2}q^3 - 2q^2 + 4\sqrt{2}q)$
$h_{14}(i, j) := h(\tilde{\varphi}_8^{i'}, \tilde{\varphi}_8^{i'-q^2 i}, \tilde{\varphi}_8^{j'}, \tilde{\varphi}_8^{j'-q^2 j})$	$i = 0, \dots, q^2 + \sqrt{2}q; j = 0, \dots, q^2 + \sqrt{2}q$ $i \neq 0, \pm j, \pm q^2 j$ $j \neq 0, \pm i, \pm q^2 i$	$\frac{1}{96}(q^4 + 2\sqrt{2}q^3 - 2q^2 - 4\sqrt{2}q)$
$h_{15}(i, j) := h(\tilde{\xi}_2^i, \tilde{\xi}_2^j, \tilde{\xi}_2^{\theta(i+j)}, \tilde{\xi}_2^{\theta(i-j)})$	$i = 0, \dots, q^2; j = 0, \dots, q^2$ $i \neq 0, \pm j, \pm(2\theta-1)j, \pm(2\theta+1)j$ $j \neq 0, \pm i, \pm(2\theta-1)i, \pm(2\theta+1)i$ $i \neq \pm(q^2+1)/3$ or $j \neq \pm(q^2+1)/3$	$\frac{q^4-10q^2+16}{48}$
$h_{16}(i) := h(\tilde{\varphi}_{12}^i, \tilde{\varphi}_{12}^{(-4\theta^3+2\theta-1)i}, \tilde{\varphi}_{12}^{(1-2\theta)i}, \tilde{\varphi}_{12}^{(4\theta^3-4\theta^2+1)i})$	$i = 0, \dots, q^4 - q^2$ $i \neq 0, \pm(q^4 - q^2 + 1)/3$	$\frac{q^4-q^2-2}{6}$
$h_{17}(i) := h(\tilde{\varphi}_{24}^{i'}, \tilde{\varphi}_{24}^{i'(q^2-q^4)i}, \tilde{\varphi}_{24}^{i'(2\theta-1)i}, \tilde{\varphi}_{24}^{i'(4\theta^3-1)i})$	$i = 0, \dots, q^4 - \sqrt{2}q^3 + q^2 - \sqrt{2}q$ $i \neq 0$	$\frac{1}{12}(q^4 - \sqrt{2}q^3 + q^2 - \sqrt{2}q)$
$h_{18}(i) := h(\tilde{\varphi}_{24}^{i'(q^4-q^2)i}, \tilde{\varphi}_{24}^{i'}, \tilde{\varphi}_{24}^{i'(2\theta+1)i}, \tilde{\varphi}_{24}^{i'(-4\theta^3-1)i})$	$i = 0, \dots, q^4 + \sqrt{2}q^3 + q^2 + \sqrt{2}q$ $i \neq 0$	$\frac{1}{12}(q^4 + \sqrt{2}q^3 + q^2 + \sqrt{2}q)$

Table B.8 (K. Shinoda [28, Tables II, V] and [17, Table A.2]) *The conjugacy classes of ${}^2F_4(q^2)$.*

Notation	Representative	$ C_2 F_4(q^2) $
$c_{1,0}$	1	$q^{24}(q^{12}+1)(q^8-1)(q^6+1)(q^2-1)$
$c_{1,1}$	$\alpha_{12}(1)$	$q^{24}(q^4+1)(q^2-1)$
$c_{1,2}$	$\alpha_{10}(1)$	$q^{20}(q^4-1)$
$c_{1,3}$	$\alpha_5(1)\alpha_{12}(1)$	$2q^{14}(q^4+1)(q^2-1)$
$c_{1,4}$	$\alpha_5(1)$	$2q^{14}(q^4+1)(q^2-1)$
$c_{1,5}$	$\alpha_7(1)\alpha_8(1)$	q^{16}
$c_{1,6}$	$\alpha_5(1)\alpha_7(1)$	q^{14}
$c_{1,7}$	$\alpha_5(1)\alpha_6(1)$	$6q^{12}$
$c_{1,8}$	$\alpha_5(1)\alpha_6(1)\alpha_8(1)$	$2q^{12}$
$c_{1,9}$	$\alpha_2(1)\alpha_6(\zeta)\alpha_8(1)$	$3q^{12}$
$c_{1,10}$	$\alpha_2(1)\alpha_4(1)$	$2q^8$
$c_{1,11}$	$\alpha_2(1)\alpha_4(1)\alpha_5(1)$	$4q^8$
$c_{1,12}$	$\alpha_2(1)\alpha_4(1)\alpha_5(1)\alpha_8(1)$	$4q^8$
$c_{1,13}$	$\alpha_2(1)\alpha_3(1)$	$2q^6$
$c_{1,14}$	$\alpha_2(1)\alpha_3(1)\alpha_5(1)$	$2q^6$
$c_{1,15}$	$\alpha_1(1)\alpha_3(1)$	$4q^4$
$c_{1,16}$	$\alpha_1(1)\alpha_2(1)\alpha_3(1)$	$4q^4$
$c_{1,17}$	$\alpha_1(1)\alpha_3(1)\alpha_5(1)$	$4q^4$
$c_{1,18}$	$\alpha_1(1)\alpha_2(1)\alpha_3(1)\alpha_5(1)$	$4q^4$
$c_{2,0}(i)$	$h_2(i)$	$q^4(q^4+1)(q^2-1)^2$
$c_{2,1}(i)$	$h_2(i)\alpha_{12}(1)$	$q^4(q^2-1)$
$c_{2,2}(i)$	$h_2(i)\alpha_5(1)$	$2q^2(q^2-1)$
$c_{2,3}(i)$	$h_2(i)\alpha_5(1)\alpha_{12}(1)$	$2q^2(q^2-1)$
$c_{3,0}(i)$	$h_3(i)$	$q^2(q^2+1)(q^2-1)^2$
$c_{3,1}(i)$	$h_3(i)\alpha_9(1)$	$q^2(q^2-1)$
$c_{4,0}(i, j)$	$h_4(i, j)$	$(q^2-1)^2$
$c_{5,0}$	h_5	$q^6(q^6+1)(q^4-1)$
$c_{5,1}$	$h_5x_{17}(1)x_{22}(1)$	$q^6(q^2+1)$
$c_{5,2}$	h_5x	$3q^4$
$c_{5,3}$	h_5x'	$3q^4$
$c_{5,4}$	h_5x''	$3q^4$
$c_{6,0}(i)$	$h_6(i)$	$q^2(q^4-1)(q^2+1)$
$c_{6,1}(i)$	$h_6(i)x_{17}(1)x_{22}(1)$	$q^2(q^2+1)$
$c_{7,0}(i)$	$h_7(i)$	q^4-1
$c_{8,0}(i)$	$h_8(i)$	$q^4(q^2-\sqrt{2}q+1)(q^4+1)(q^2-1)$
$c_{8,1}(i)$	$h_8(i)x_{21}(1)x_{24}(1)$	$q^4(q^2-\sqrt{2}q+1)$
$c_{8,2}(i)$	$h_8(i)x_8(1)x_{16}(1)x_{21}(1)$	$2q^2(q^2-\sqrt{2}q+1)$
$c_{8,3}(i)$	$h_8(i)x_8(1)x_{16}(1)x_{24}(1)$	$2q^2(q^2-\sqrt{2}q+1)$
$c_{9,0}(i)$	$h_9(i)$	$(q^2-\sqrt{2}q+1)(q^2-1)$
$c_{10,0}(i)$	$h_{10}(i)$	$q^4(q^2+\sqrt{2}q+1)(q^4+1)(q^2-1)$
$c_{10,1}(i)$	$h_{10}(i)x_{21}(1)x_{24}(1)$	$q^4(q^2+\sqrt{2}q+1)$
$c_{10,2}(i)$	$h_{10}(i)x_8(1)x_{16}(1)x_{21}(1)$	$2q^2(q^2+\sqrt{2}q+1)$
$c_{10,3}(i)$	$h_{10}(i)x_8(1)x_{16}(1)x_{24}(1)$	$2q^2(q^2+\sqrt{2}q+1)$
$c_{11,0}(i)$	$h_{11}(i)$	$(q^2+\sqrt{2}q+1)(q^2-1)$
$c_{12,0}(i, j)$	$h_{12}(i, j)$	q^4+1
$c_{13,0}(i, j)$	$h_{13}(i, j)$	$(q^2-\sqrt{2}q+1)^2$
$c_{14,0}(i, j)$	$h_{14}(i, j)$	$(q^2+\sqrt{2}q+1)^2$
$c_{15,0}(i, j)$	$h_{15}(i, j)$	$(q^2+1)^2$
$c_{16,0}(i)$	$h_{16}(i)$	q^4-q^2+1
$c_{17,0}(i)$	$h_{17}(i)$	$q^4-\sqrt{2}q^3+q^2-\sqrt{2}q+1$
$c_{18,0}(i)$	$h_{18}(i)$	$q^4+\sqrt{2}q^3+q^2+\sqrt{2}q+1$

Table B.9 *Parameterization of the types of Lusztig series of ${}^2F_4(q^2)$.*

No.	Dynkin type	Π	w
g_1	2F_4	$[r_1^\vee, r_2^\vee, r_3^\vee, r_4^\vee]$	1
g_2	2B_2	$[r_8^\vee, r_{16}^\vee]$	1
g_3	A_1	$[r_{15}^\vee, r_{20}^\vee]$	1
g_4	A_0	$[\]$	1
g_5	2A_2	if $\theta \equiv 1 \pmod 3$: $[r_6^\vee, r_{11}^\vee, r_{12}^\vee, r_{16}^\vee]$ if $\theta \equiv -1 \pmod 3$: $[r_8^\vee, r_9^\vee, r_{10}^\vee, r_{18}^\vee]$	w_{r_1}
g_6	A_1	$[r_{17}^\vee, r_{22}^\vee]$	w_{r_1}
g_7	A_0	$[\]$	w_{r_1}
g_8	2B_2	$[r_8^\vee, r_{16}^\vee]$	$w_{r_1} w_{r_2} w_{r_1} w_{r_3} w_{r_2} w_{r_1} w_{r_3} w_{r_2}$
g_9	A_0	$[\]$	w_{r_3}
g_{10}	2B_2	$[r_8^\vee, r_{16}^\vee]$	$w_{r_1} w_{r_2} w_{r_1} w_{r_3} w_{r_2} w_{r_1} w_{r_3} w_{r_2}$
g_{11}	A_0	$[\]$	$w_{r_2} w_{r_3} w_{r_2}$
g_{12}	A_0	$[\]$	$w_{r_1} w_{r_2} w_{r_1} w_{r_3} w_{r_2} w_{r_1} w_{r_3} w_{r_2}$
g_{13}	A_0	$[\]$	$w_{r_1} w_{r_2} w_{r_3} w_{r_2} w_{r_1} w_{r_3}$
g_{14}	A_0	$[\]$	$w_{r_2} w_{r_3} w_{r_2} w_{r_4} (w_{r_3} w_{r_2} w_{r_1} w_{r_3} w_{r_2} w_{r_3} w_{r_4})^2$
g_{15}	A_0	$[\]$	$w_{r_1} w_{r_2} w_{r_3} w_{r_4} w_{r_3} w_{r_2} w_{r_1} w_{r_3} w_{r_2} w_{r_4} w_{r_3} w_{r_2}$
g_{16}	A_0	$[\]$	$w_{r_2} w_{r_3} w_{r_2} w_{r_1}$
g_{17}	A_0	$[\]$	$w_{r_1} w_{r_2}$
g_{18}	A_0	$[\]$	$w_{r_1} w_{r_4} w_{r_3} w_{r_2} w_{r_1} w_{r_3} w_{r_2} w_{r_4} w_{r_3} w_{r_2}$

Table B.10 *Representatives for the semisimple conjugacy classes of $\mathbf{G}^{*F^*}$.*

Representative	Parameters	Number of Classes
$g_1 := s(0, 0, 0, 0)$		1
$g_2(k) := s\left(0, \frac{(2\theta+2)k}{q^2-1}, \frac{(4\theta+2)k}{q^2-1}, 0\right)$	$k = 0, \dots, q^2 - 2$ $k \neq 0$	$\frac{q^2-2}{2}$
$g_3(k) := s\left(\frac{2\theta k}{q^2-1}, \frac{(4\theta+2)k}{q^2-1}, \frac{(4\theta+4)k}{q^2-1}, \frac{2k}{q^2-1}\right)$	$k = 0, \dots, q^2 - 2$ $k \neq 0$	$\frac{q^2-2}{2}$
$g_4(k, l) := s_1(k, l)$	$k = 0, \dots, q^2 - 2; l = 0, \dots, q^2 - 2$ $k \neq 0, \pm l, \pm(2\theta - 1)l$ $l \neq 0, \pm k, \pm(2\theta - 1)k$	$\frac{q^4-10q^2+16}{16}$
$g_5 := s\left(\frac{1}{3}, 0, 0, \frac{\theta}{3}\right)$		1
$g_6(k) := s\left(\frac{\theta^2 k}{q^2+1}, 0, 0, \frac{\theta k}{q^2+1}\right)$	$k = 0, \dots, q^2$ $k \neq 0, \frac{q^2+1}{3}, \frac{2(q^2+1)}{3}$	$\frac{q^2-2}{2}$
$g_7(k) := s_2(k)$	$k = 0, \dots, q^4 - 2$ $k \neq (q^2 - 1)m, m = 0, \dots, q^2$ $k \neq (q^2 + 1)m, m = 0, \dots, q^2 - 2$	$\frac{q^4-2q^2}{4}$
$g_8(k) := s\left(0, \frac{(\theta-1)k}{q^2-\sqrt{2}q+1}, \frac{-k}{q^2-\sqrt{2}q+1}, 0\right)$	$k = 0, \dots, q^2 - \sqrt{2}q$ $k \neq 0$	$\frac{q^2-\sqrt{2}q}{4}$
$g_9(k) := s_3(k)$	$k = 0, \dots, q^4 - \sqrt{2}q^3 + \sqrt{2}q - 2$ $k \neq (q^2 - 1)m, m = 0, \dots, q^2 - \sqrt{2}q$ $k \neq (q^2 - \sqrt{2}q + 1)m, m = 0, \dots, q^2 - 2$	$\frac{1}{8}(q^4 - \sqrt{2}q^3 - 2q^2 + 2\sqrt{2}q)$
$g_{10}(k) := s\left(0, \frac{(-\theta-1)k}{q^2+\sqrt{2}q+1}, \frac{-k}{q^2+\sqrt{2}q+1}, 0\right)$	$k = 0, \dots, q^2 + \sqrt{2}q$ $k \neq 0$	$\frac{q^2+\sqrt{2}q}{4}$
$g_{11}(k) := s_4(k)$	$k = 0, \dots, q^4 + \sqrt{2}q^3 - \sqrt{2}q - 2$ $k \neq (q^2 - 1)m, m = 0, \dots, q^2 + \sqrt{2}q$ $k \neq (q^2 + \sqrt{2}q + 1)m, m = 0, \dots, q^2 - 2$	$\frac{1}{8}(q^4 + \sqrt{2}q^3 - 2q^2 - 2\sqrt{2}q)$
$g_{12}(k, l) := s_5(k)$	$k = 0, \dots, q^2 - \sqrt{2}q; l = 0, \dots, q^2 + \sqrt{2}q$ $k, l \neq 0$	$\frac{q^2(q^2-2)}{16}$
$g_{13}(k, l) := s_6(k, l)$	$k = 0, \dots, q^2 - \sqrt{2}q; b = l, \dots, q^2 - \sqrt{2}q$ $k \neq 0, \pm l, \pm q^2 l$ $l \neq 0, \pm k, \pm q^2 k$	$\frac{1}{96}(q^4 - 2\sqrt{2}q^3 - 2q^2 + 4\sqrt{2}q)$
$g_{14}(k, l) := s_7(k, l)$	$k = 0, \dots, q^2 + \sqrt{2}q; l = 0, \dots, q^2 + \sqrt{2}q$ $k \neq 0, \pm l, \pm q^2 l$ $l \neq 0, \pm k, \pm q^2 k$	$\frac{1}{96}(q^4 + 2\sqrt{2}q^3 - 2q^2 - 4\sqrt{2}q)$
$g_{15}(k, l) := s_8(k, l)$	$k = 0, \dots, q^2; l = 0, \dots, q^2$ $k \neq 0, \pm l, \pm(2\theta - 1)l, \pm(2\theta + 1)l$ $l \neq 0, \pm k, \pm(2\theta - 1)k, \pm(2\theta + 1)k$ $k \neq \pm(q^2 + 1)/3$ or $l \neq \pm(q^2 + 1)/3$	$\frac{q^4-10q^2+16}{48}$
$g_{16}(k) := s_9(k)$	$k = 0, \dots, q^4 - q^2$ $k \neq 0, \pm(q^4 - q^2 + 1)/3$	$\frac{q^4-q^2-2}{6}$
$g_{17}(k) := s_{10}(k)$	$k = 0, \dots, q^4 - \sqrt{2}q^3 + q^2 - \sqrt{2}q$ $k \neq 0$	$\frac{1}{12}(q^4 - \sqrt{2}q^3 + q^2 - \sqrt{2}q)$
$g_{18}(k) := s_{11}(k)$	$k = 0, \dots, q^4 + \sqrt{2}q^3 + q^2 + \sqrt{2}q$ $k \neq 0$	$\frac{1}{12}(q^4 + \sqrt{2}q^3 + q^2 + \sqrt{2}q)$

Table B.11 Values of the class functions $\pm f_8(k)$ and $\pm f_{10}(k)$ in the Lusztig series of types g_8, g_{10} . All values not listed here are zero.

	$c_{1,3}$		$c_{1,4}$	
$f_8(k)$	$\sqrt{2}q^5\varepsilon_4(q^2 + \sqrt{2}q + 1)(q^2 - 1)$		$-\sqrt{2}q^5\varepsilon_4(q^2 + \sqrt{2}q + 1)(q^2 - 1)$	
$f_{10}(k)$	$\sqrt{2}q^5\varepsilon_4(q^2 - \sqrt{2}q + 1)(q^2 - 1)$		$-\sqrt{2}q^5\varepsilon_4(q^2 - \sqrt{2}q + 1)(q^2 - 1)$	
	$c_{1,11}$		$c_{1,12}$	
$f_8(k)$	$-\sqrt{2}q^2\varepsilon_4(2q + \sqrt{2})$		$\sqrt{2}q^2\varepsilon_4(2q + \sqrt{2})$	
$f_{10}(k)$	$\sqrt{2}q^2\varepsilon_4(2q - \sqrt{2})$		$-\sqrt{2}q^2\varepsilon_4(2q - \sqrt{2})$	
	$c_{1,15}$	$c_{1,16}$	$c_{1,17}$	$c_{1,18}$
$f_8(k)$	$-\sqrt{2}q\varepsilon_4$	$\sqrt{2}q\varepsilon_4$	$-\sqrt{2}q\varepsilon_4$	$\sqrt{2}q\varepsilon_4$
$f_{10}(k)$	$-\sqrt{2}q\varepsilon_4$	$\sqrt{2}q\varepsilon_4$	$-\sqrt{2}q\varepsilon_4$	$\sqrt{2}q\varepsilon_4$
	$c_{8,2}(i)$			
$f_8(k)$	$\sqrt{2}q\varepsilon_4(\varphi_8''^{qiq/\sqrt{2}} + \varphi_8''^{-qiq/\sqrt{2}} + \varphi_8''^{(q/\sqrt{2}-1)ik} + \varphi_8''^{-(q/\sqrt{2}-1)ik})$			
$f_{10}(k)$	0			
	$c_{8,3}(i)$			
$f_8(k)$	$-\sqrt{2}q\varepsilon_4(\varphi_8''^{qiq/\sqrt{2}} + \varphi_8''^{-qiq/\sqrt{2}} + \varphi_8''^{(q/\sqrt{2}-1)ik} + \varphi_8''^{-(q/\sqrt{2}-1)ik})$			
$f_{10}(k)$	0			
	$c_{10,2}(i)$			
$f_8(k)$	0			
$f_{10}(k)$	$\sqrt{2}q\varepsilon_4(\varphi_8'^{qiq/\sqrt{2}} + \varphi_8'^{-qiq/\sqrt{2}} + \varphi_8'^{(q/\sqrt{2}+1)ik} + \varphi_8'^{-(q/\sqrt{2}+1)ik})$			
	$c_{10,3}(i)$			
$f_8(k)$	0			
$f_{10}(k)$	$-\sqrt{2}q\varepsilon_4(\varphi_8'^{qiq/\sqrt{2}} + \varphi_8'^{-qiq/\sqrt{2}} + \varphi_8'^{(q/\sqrt{2}+1)ik} + \varphi_8'^{-(q/\sqrt{2}+1)ik})$			

Table B.12 Values of the irreducible characters $G\chi_{42}(k), \dots, G\chi_{49}(k)$ of the group $G := {}^2F_4(q^2)$ on some conjugacy classes.

	$c_{1,0}$
$G\chi_{42}(k)$	$(q-1)(q+1)(q^2+1)^2(q^4-q^2+1)(q^8-q^4+1)(q^2+\sqrt{2}q+1)$
$G\chi_{43}(k)$	$\frac{q}{\sqrt{2}}(q-1)^2(q+1)^2(q^2+1)^2(q^4-q^2+1)(q^8-q^4+1)(q^2+\sqrt{2}q+1)$
$G\chi_{44}(k)$	$\frac{q}{\sqrt{2}}(q-1)^2(q+1)^2(q^2+1)^2(q^4-q^2+1)(q^8-q^4+1)(q^2+\sqrt{2}q+1)$
$G\chi_{45}(k)$	$q^4(q-1)(q+1)(q^2+1)^2(q^4-q^2+1)(q^8-q^4+1)(q^2+\sqrt{2}q+1)$
$G\chi_{46}(k)$	$(q-1)(q+1)(q^2+1)^2(q^4-q^2+1)(q^8-q^4+1)(q^2-\sqrt{2}q+1)$
$G\chi_{47}(k)$	$\frac{q}{\sqrt{2}}(q-1)^2(q+1)^2(q^2+1)^2(q^4-q^2+1)(q^8-q^4+1)(q^2-\sqrt{2}q+1)$
$G\chi_{48}(k)$	$\frac{q}{\sqrt{2}}(q-1)^2(q+1)^2(q^2+1)^2(q^4-q^2+1)(q^8-q^4+1)(q^2-\sqrt{2}q+1)$
$G\chi_{49}(k)$	$q^4(q-1)(q+1)(q^2+1)^2(q^4-q^2+1)(q^8-q^4+1)(q^2-\sqrt{2}q+1)$

	$c_{1,11}$
$G\chi_{42}(k)$	$-(q^2+\sqrt{2}q+1)$
$G\chi_{43}(k)$	$q^2+\frac{1}{2}\sqrt{2}q-\varepsilon_4\sqrt{2}q^3-\varepsilon_4q^2$
$G\chi_{44}(k)$	$q^2+\frac{1}{2}\sqrt{2}q+\varepsilon_4\sqrt{2}q^3+q^2\varepsilon_4$
$G\chi_{45}(k)$	0
$G\chi_{46}(k)$	$-(q^2-\sqrt{2}q+1)$
$G\chi_{47}(k)$	$-q^2+\frac{1}{2}\sqrt{2}q+\varepsilon_4\sqrt{2}q^3-\varepsilon_4q^2$
$G\chi_{48}(k)$	$-q^2+\frac{1}{2}\sqrt{2}q-\varepsilon_4\sqrt{2}q^3+q^2\varepsilon_4$
$G\chi_{49}(k)$	0

	$c_{8,2}(i)$
$G\chi_{42}(k)$	$-\varphi_8''^{\frac{1}{2}iq\sqrt{2}k-ik}-\varphi_8''^{\sqrt{2}qik-ik}-\varphi_8''^{-\sqrt{2}qik+ik}-\varphi_8''^{-\frac{1}{2}iq\sqrt{2}k+ik}$ $-\varphi_8''^{\frac{1}{2}iq\sqrt{2}k}-\varphi_8''^{-\frac{1}{2}iq\sqrt{2}k}-\varphi_8''^{-ik}-\varphi_8''^{ik}-1$
$G\chi_{43}(k)$	$\frac{1}{2}q\sqrt{2}\varepsilon_4\varphi_8''^{\frac{1}{2}iq\sqrt{2}k}+\frac{1}{2}q\sqrt{2}\varepsilon_4\varphi_8''^{-\frac{1}{2}iq\sqrt{2}k+ik}$ $+\frac{1}{2}q\sqrt{2}\varepsilon_4\varphi_8''^{\frac{1}{2}iq\sqrt{2}k-ik}+\frac{1}{2}q\sqrt{2}\varepsilon_4\varphi_8''^{-\frac{1}{2}iq\sqrt{2}k}$ $+\varphi_8''^{\frac{1}{2}iq\sqrt{2}k-ik}+\varphi_8''^{-\frac{1}{2}iq\sqrt{2}k+ik}+\varphi_8''^{\frac{1}{2}iq\sqrt{2}k}+\varphi_8''^{-\frac{1}{2}iq\sqrt{2}k}+1$
$G\chi_{44}(k)$	$-\frac{1}{2}q\sqrt{2}\varepsilon_4\varphi_8''^{\frac{1}{2}iq\sqrt{2}k}-\frac{1}{2}q\sqrt{2}\varepsilon_4\varphi_8''^{-\frac{1}{2}iq\sqrt{2}k+ik}$ $-\frac{1}{2}q\sqrt{2}\varepsilon_4\varphi_8''^{\frac{1}{2}iq\sqrt{2}k-ik}-\frac{1}{2}q\sqrt{2}\varepsilon_4\varphi_8''^{-\frac{1}{2}iq\sqrt{2}k}+\varphi_8''^{\frac{1}{2}iq\sqrt{2}k-ik}$ $+\varphi_8''^{-\frac{1}{2}iq\sqrt{2}k+ik}+\varphi_8''^{\frac{1}{2}iq\sqrt{2}k}+\varphi_8''^{-\frac{1}{2}iq\sqrt{2}k}+1$
$G\chi_{45}(k)$	$1+\varphi_8''^{\frac{1}{2}iq\sqrt{2}k-ik}+\varphi_8''^{-\frac{1}{2}iq\sqrt{2}k+ik}+\varphi_8''^{\frac{1}{2}iq\sqrt{2}k}+\varphi_8''^{-\frac{1}{2}iq\sqrt{2}k}$
$G\chi_{46}(k)$	-1
$G\chi_{47}(k)$	1
$G\chi_{48}(k)$	1
$G\chi_{49}(k)$	1

Table B.12 (cont.)

	$c_{10,2}(i)$
$G\chi_{42}(k)$	-1
$G\chi_{43}(k)$	-1
$G\chi_{44}(k)$	-1
$G\chi_{45}(k)$	1
$G\chi_{46}(k)$	$-\varphi_8'^{-\sqrt{2}qik-ik} - \varphi_8'^{-\frac{1}{2}iq\sqrt{2}k-ik} - \varphi_8'^{\frac{1}{2}iq\sqrt{2}k+ik} - \varphi_8'^{\sqrt{2}qik+ik}$ $-\varphi_8'^{\frac{1}{2}iq\sqrt{2}k} - \varphi_8'^{-\frac{1}{2}iq\sqrt{2}k} - \varphi_8'^{-ik} - \varphi_8'^{ik} - 1$
$G\chi_{47}(k)$	$\frac{1}{2}q\sqrt{2}\varepsilon_4\varphi_8'^{\frac{1}{2}iq\sqrt{2}k} + \frac{1}{2}q\sqrt{2}\varepsilon_4\varphi_8'^{-\frac{1}{2}iq\sqrt{2}k-ik}$ $+ \frac{1}{2}q\sqrt{2}\varepsilon_4\varphi_8'^{-\frac{1}{2}iq\sqrt{2}k} + \frac{1}{2}q\sqrt{2}\varepsilon_4\varphi_8'^{\frac{1}{2}iq\sqrt{2}k+ik} - \varphi_8'^{-\frac{1}{2}iq\sqrt{2}k-ik}$ $-\varphi_8'^{\frac{1}{2}iq\sqrt{2}k+ik} - \varphi_8'^{-\frac{1}{2}iq\sqrt{2}k} - \varphi_8'^{\frac{1}{2}iq\sqrt{2}k} - 1$
$G\chi_{48}(k)$	$-\frac{1}{2}q\sqrt{2}\varepsilon_4\varphi_8'^{\frac{1}{2}iq\sqrt{2}k} - \frac{1}{2}q\sqrt{2}\varepsilon_4\varphi_8'^{-\frac{1}{2}iq\sqrt{2}k-ik}$ $-\frac{1}{2}q\sqrt{2}\varepsilon_4\varphi_8'^{-\frac{1}{2}iq\sqrt{2}k} - \frac{1}{2}q\sqrt{2}\varepsilon_4\varphi_8'^{\frac{1}{2}iq\sqrt{2}k+ik} - \varphi_8'^{-\frac{1}{2}iq\sqrt{2}k-ik}$ $-\varphi_8'^{\frac{1}{2}iq\sqrt{2}k+ik} - \varphi_8'^{-\frac{1}{2}iq\sqrt{2}k} - \varphi_8'^{\frac{1}{2}iq\sqrt{2}k} - 1$
$G\chi_{49}(k)$	$1 + \varphi_8'^{\frac{1}{2}iq\sqrt{2}k} + \varphi_8'^{-\frac{1}{2}iq\sqrt{2}k-ik} + \varphi_8'^{-\frac{1}{2}iq\sqrt{2}k} + \varphi_8'^{\frac{1}{2}iq\sqrt{2}k+ik}$

REFERENCES

- [1] J. L. ALPERIN, The main problem of block theory, in Proceedings of the Conference on Finite Groups, Univ. Utah, Park City, UT, Academic Press, New York, 1975 pp. 341–356. [1](#)
- [2] J. L. ALPERIN, Weights for finite groups, in The Arcata Conference on Representations of Finite Groups, Proc. Sympos. Pure Math, **47**, AMS, Providence, RI, 1987 pp. 369–379. [1](#)
- [3] J. AN, The Alperin and Dade conjectures for the Ree groups ${}^2F_4(q^2)$ in non-defining characteristics, *J. Algebra*, **203** (1998), 30–49. [1](#)
- [4] J. AN, Dade’s conjecture for the Tits group, *New Zealand Journal of Mathematics*, **25** (1996), 107–131. [1](#)
- [5] J. AN, F. HIMSTEDT AND S. HUANG, Uno’s invariant conjecture for Steinberg’s triality groups in defining characteristic, *J. Algebra*, **316** (2007), 79–108. [3](#)
- [6] N. BURGOS AND C. WILLIAMSON, On a theorem of Borel and Tits for finite Chevalley groups, *Arch. Math. (Basel)*, **27** (1976), 489–491. [1](#), [11](#)
- [7] R. W. CARTER, Finite groups of Lie type - Conjugacy classes and complex characters, A Wiley-Interscience publication, Chichester, 1985. [27](#), [28](#), [31](#)
- [8] B. CHAR, K. GEDDES, G. GONNET, B. LEONG, M. MONAGAN AND S. WATT, Maple V, Language Reference Manual, Springer, 1991. [1](#)
- [9] E. C. DADE, Counting characters in blocks I, *Invent. Math.*, **109** (1992), 187–210. [1](#)
- [10] E. C. DADE, Counting characters in blocks II, *J. reine angew. Math.*, **448** (1994), 97–190. [1](#)
- [11] E. C. DADE, Counting characters in blocks 2.9, in R. SOLOMON, ed., Representation Theory of Finite Groups, 1997 pp. 45–59. [1](#), [3](#)
- [12] F. DIGNE AND J. MICHEL, Representations of finite groups of Lie type, Cambridge University Press, Cambridge, 1990. [28](#), [33](#)
- [13] The GAP Group, GAP – Groups, Algorithms, and Programming, Version 4.4.12 (2008), <http://www.gap-system.org>. [27](#)
- [14] M. GECK, G. HISS, F. LÜBECK, G. MALLE AND G. PFEIFFER, CHEVIE – A system for computing and processing generic character tables for finite groups of Lie type, Weyl groups and Hecke algebras, *Appl. Algebra Engrg. Comm. Comput.*, **7** (1996), 175–210. [1](#), [5](#), [27](#)
- [15] F. HIMSTEDT, On the decomposition numbers of the Ree groups ${}^2F_4(q^2)$ in non-defining characteristic, submitted. [2](#), [27](#)
- [16] F. HIMSTEDT AND S. HUANG, Dade’s invariant conjecture for Steinberg’s triality groups ${}^3D_4(2^n)$ in defining characteristic, *J. Algebra*, **316** (2007), 802–827. [1](#)

- [17] F. HIMSTEDT AND S. HUANG, Character table of a Borel subgroup of the Ree groups ${}^2F_4(q^2)$, *LMS J. Comput. Math.*, **12** (2009), 1–53. [1](#), [5](#), [6](#), [11](#), [12](#), [14](#), [27](#), [28](#), [29](#), [32](#), [40](#), [41](#)
- [18] F. HIMSTEDT AND S. HUANG, Character tables of the maximal parabolic subgroups of the Ree groups ${}^2F_4(q^2)$, to appear in *LMS J. Comput. Math.* [1](#), [5](#), [6](#), [11](#), [12](#)
- [19] J. HUMPHREYS, Defect groups for finite groups of Lie type, *Math. Z.*, **119** (1971), 149–152. [11](#)
- [20] I. M. ISAACS AND G. NAVARRO, New refinements of the McKay conjecture for arbitrary finite groups, *Ann. of Math.*, **156** (2002), 333–344. [1](#)
- [21] I. M. ISAACS, G. MALLE AND G. NAVARRO, A reduction theorem for the McKay conjecture, *Invent. Math.*, **170** (2007), 33–101. [1](#)
- [22] R. KNÖRR AND G. ROBINSON, Some remarks on a conjecture of Alperin, *J. London Math. Soc. (2)*, **39** (1989), 48–60. [1](#)
- [23] C. KÖHLER, Unipotente Charaktere und Zerlegungszahlen der endlichen Chevalleygruppen vom Typ F_4 , Dissertation, RWTH Aachen, Germany, 2006. [29](#)
- [24] F. LÜBECK, Charaktertafeln für die Gruppen $\mathrm{CSp}_6(q)$ mit ungeradem q und $\mathrm{Sp}_6(q)$ mit geradem q , Dissertation, Universität Heidelberg, Germany, 1993. [27](#), [28](#), [29](#), [30](#), [31](#), [32](#)
- [25] G. MALLE, Die unipotenten Charaktere von ${}^2F_4(q^2)$, *Comm. Alg.* **18** (1990), 2361–2381. [27](#)
- [26] J. MCKAY, A new invariant for simple groups, *Notices Amer. Math. Soc.*, **18** (1971), 397. [1](#)
- [27] R. REE, A family of simple groups associated with the simple Lie algebra of type (F_4) , *Amer. J. Math.* **83** (1961), 401–420. [27](#)
- [28] K. SHINODA, The conjugacy classes of the finite Ree groups of type (F_4) , *J. Fac. Sci. Univ. Tokyo Sect. I A Math.*, **22** (1975), 1–15. [27](#), [28](#), [29](#), [33](#), [35](#), [36](#), [40](#), [41](#)
- [29] K. UNO, Conjectures on character degrees for the simple Thompson group, *Osaka J. Math.*, **41** (2004), 11–36. [1](#)

F.H.: TECHNISCHE UNIVERSITÄT MÜNCHEN, ZENTRUM MATHEMATIK – M11, BOLTZMANNSTR. 3, 85748 GARCHING, GERMANY

S.H.: DEPARTMENT OF MATHEMATICS, NATIONAL CHENG KUNG UNIVERSITY, NO. 1 DASYUE RD, TAINAN CITY 70101, TAIWAN

E-mail address: F.H.: himstedt@ma.tum.de

E-mail address: S.H.: shua3@mail.ncku.edu.tw